

08/21/97

PATENT APPLICATION TRANSMITTAL LETTER

PATENT GROUP ATTY DOCKET SUV-003.04  
 FOLEY, HOAG & ELIOT LLP  
 ONE POST OFFICE SQUARE  
 BOSTON, MA 02109-2170

To the Assistant Commissioner for Patents:

Transmitted herewith for filing under 35 U.S.C. 111 and 37 CFR 1.53 is the patent application of:

MATTHEW P. SCOTT, LISA V. GOODRICH, RONALD L. JOHNSON, and ERVIN EPSTEIN JR.

entitled Patched Genes and Uses Related Thereto

Enclosed are:

- (X) 80 pages of written description, claims and abstract.  
 (X) 5 sheets of drawings.  
 ( ) an assignment of the invention to \_\_\_\_\_  
 ( ) executed declaration of the inventors.  
 ( ) a certified copy of a \_\_\_\_\_ application.  
 ( ) a verified statement to establish small entity status under 37 CFR 1.9 and 1.27.  
 (X) other: unexecuted Declaration, Petition and Power of Attorney.

CLAIMS AS FILED

	# FILED	# EXTRA	Rate	FEE	Rate (Small Entity)	FEE
BASIC FEE			\$770		\$385	
TOTAL CLAIMS	-20 =	0	x \$22		x \$11	
INDEPENDENT CLAIMS	-3 =	0	x \$80		x \$40	
MULTIPLE DEPENDENT CLAIMS			\$260		\$130	
* NUMBER EXTRA MUST BE ZERO OR LARGER			TOTAL			\$

- ( ) A check in the amount of \$ \_\_\_\_\_ to cover the filing fee is enclosed.  
 ( ) The Commissioner is hereby authorized to charge and credit Deposit Account No. 06-1448 as described below. A duplicate copy of this sheet is enclosed.  
 ( ) Charge the amount of \$ \_\_\_\_\_ as filing fee.  
 ( ) Credit any overpayment.  
 ( ) Charge any additional filing fees required under 37 CFR 1.16 and 1.17.

Certificate of Express Mail

Express Mail Label: EM 528 936 719 US

I hereby certify that this correspondence is being deposited with the United States Postal Service as Express Mail, postage prepaid, "Post Office to Addressee", in an envelope addressed to the Assistant Commissioner for Patents, Box Patent Application, Washington, D.C. 20231 on this date of August 21, 1997.

8-21-97

Date of Deposit

Ariel I. Collazo

Ariel I. Collazo

Matthew P. Vincent, Ph.D.  
 Reg. No. 36,709  
 Attorney for Applicants

## Patched Genes and Uses Related Thereto

---

### Related Applications

This application is a continuation-in-part of U.S.S.N. 08/656,055, which is a continuation-in-  
5 part of U.S.S.N. 08/540,406, which is a continuation-in-part of U.S.S.N. 08/317,745 (now  
abandoned). The specifications of each of these prior applications are incorporated herein by  
reference.

### Background of the Invention

10 Segment polarity genes were originally discovered as mutations in flies that change  
the pattern of body segment structures. Mutations in these genes cause animals to develop  
changed patterns on the surfaces of body segments; the changes affecting the pattern along  
the head to tail axis. Among the genes in this class are *hedgehog*, which encodes a secreted  
protein (HH), and *patched*, which encodes a protein structurally similar to transporter  
15 proteins, having twelve transmembrane domains (*ptc*), with two conserved glycosylation  
signals.

The *hedgehog* gene of flies has at least three vertebrate relatives- *Sonic hedgehog*  
(*Shh*); *Indian hedgehog* (*Ihh*), and *Desert hedgehog* (*Dhh*). *Shh* is expressed in a group of  
cells, at the posterior of each developing limb bud, that have an important role in signaling  
20 polarity to the developing limb. The *Shh* protein product, SHH, is a critical trigger of  
posterior limb development, and is also involved in polarizing the neural tube and somites  
along the dorsal ventral axis. Based on genetic experiments in flies, *patched* and *hedgehog*  
have antagonistic effects in development. The *patched* gene product, *ptc*, is widely expressed  
in fetal and adult tissues, and plays an important role in regulation of development. *Ptc*  
25 downregulates transcription of itself, members of the transforming growth factor and *Wnt*  
gene families, and possibly other genes. Among other activities, HH upregulates expression  
of *patched* and other genes that are negatively regulated by *patched*.

It is of interest that many genes involved in the regulation of growth and control of  
cellular signaling are also involved in oncogenesis. Such genes may be oncogenes, which are  
30 typically upregulated in tumor cells, or tumor suppressor genes, which are down-regulated or  
absent in tumor cells. Malignancies may arise when a tumor suppressor is lost and/or an  
oncogene is inappropriately activated. Familial predisposition to cancer may occur when  
there is a mutation, such as loss of an allele encoding a suppressor gene, present in the  
germline DNA of an individual.

The most common form of cancer in the United States is basal cell carcinoma of the skin. While sporadic cases are very common, there are also familial syndromes, such as the basal cell nevus syndrome (BCNS). The familial syndrome has many features indicative of abnormal embryonic development, indicating that the mutated gene also plays an important role in development of the embryo. A loss of heterozygosity of chromosome 9q alleles in both familial and sporadic carcinomas suggests that a tumor suppressor gene is present in this region. The high incidence of skin cancer makes the identification of this putative tumor suppressor gene of great interest for diagnosis, therapy, and drug screening.

## Relevant Literature

Descriptions of *patched*, by itself or its role with *hedgehog* may be found in Hooper and Scott (1989) *Cell* **59**:751-765; and Nakano *et al.* (1989) *Nature* **341** :508-513. Both of these references also describe the sequence for *Drosophila patched*. Discussions of the role of *hedgehog* include Riddle *et al.* (1993) *Cell* **75**:1401-1416-, Echelard *et al.* (1993) *Cell* **75**:1417-1430- Krauss *et al.* (1993) *Cell* **75**:1431-1444 (1993); Tabata and Kornberg (1994) *Cell* **76**:89-102; Heemskerk and DiNardo (1994) *Cell* **76**:449-460; and Roelink *et al.* (1994) *Cell* **76**:761-775.

Mapping of deleted regions on chromosome 9 in skin cancers is described in Habuchi *et al.* (1995) *Oncogene* **11**: 1 671-1674, Quinn *et al.* (1 994) *Genes Chromosome Cancer* **11**:222-225; Quinn *et al.* (1994) *J. Invest. Dermatol.* **102**:300-303; and Wicking *et al.* (1994) *Genomics* **22**:505-51 1.

Gorlin (1987) *Medicine* **66**:98-113 reviews nevoid basal cell carcinoma syndrome. The syndrome shows autosomal dominant inheritance with probably complete penetrance. About 60% of the cases represent new mutations. Developmental abnormalities found with this syndrome include rib and craniofacial abnormalities, polydactyly, syndactyly and spina bifida. Tumors found with the syndrome include basal cell carcinomas, fibromas of the ovaries and heart, cysts of the skin, jaws and mesentery, meningiomas and medulloblastomas.

## Summary of the Invention

Isolated nucleotide compositions and sequences are provided for *patched* (*ptc*) genes, including mammalian, e.g. human and mouse, and invertebrate homologs. Decreased expression of *ptc* is associated with the occurrence of human cancers, particularly basal cell carcinomas and other tumors of epithelial tissues such as the skin. The cancers may be familial, having as a component of risk a germline mutation in the gene, or may be sporadic. *Ptc*, and its antagonist *hedgehog*, are useful in creating transgenic animal models for these human cancers. The *ptc* nucleic acid compositions find use in identifying homologous or

related genes; in producing compositions that modulate the expression or function of its encoded protein, *ptc*; for gene therapy; mapping functional regions of the protein- and in studying associated physiological pathways. In addition, modulation of the gene activity *in vivo* is used for prophylactic and therapeutic purposes, such as treatment of cancer, identification of cell type based on expression, and the like. *Ptc*, anti-*ptc* antibodies and *ptc* nucleic acid sequences are useful as diagnostics for a genetic predisposition to cancer or developmental abnormality syndromes, and to identify specific cancers having mutations in this gene.

### Brief Description of the Drawings

Fig. 1 is a graph having a restriction map of about 10 kbp of the 5' region upstream from the initiation codon of *Drosophila patched* gene and bar graphs of constructs of truncated portions of the 5' region joined to fl-galactosidase, where the constructs are introduced into fly cell lines for the production of embryos. The expression of fl-gal in the embryos is indicated in the right-hand table during early and late development of the embryo. The greater the number of +'s, the more intense the staining.

Fig. 2 shows a summary of mutations found in the human *patched* gene locus that are associated with basal cell nevus syndrome. Mutation (1) is found in sporadic basal cell carcinoma, and is a C to T transition in exon 3 at nucleotide 523 of the coding sequence, changing Leu 175 to Phe in the first extracellular loop. Mutations 2-4 are found in hereditary basal carcinoma nevus syndrome. (2) is an insertion of 9 bp at nucleotide 2445, resulting in the insertion of an additional 3 amino acids after amino acid 815. (3) is a deletion of 11 bp, which removes nt 2442-2452 from the coding sequence. The resulting frameshift truncates the open reading frame after amino acid 813, 'just after the seventh transmembrane domain. (4) is a G to C alteration that changes two conserved nucleotides of the 3' splice site adjacent to exon 10, creating a non-functional splice site that truncates the protein after amino acid 449, in the second transmembrane region.

Fig. 3 (panels A-B) illustrates the generation of *ptc* mutations. (A) The *ptc* mutant allele was generated by homologous recombination between the KO1 targeting vector and *ptc*. External probe A detected a 3' EcoRV polymorphism on blots and probe B detected a 5' SacI polymorphism. Exons are numbered. (B) Transmission of the *ptc*<sup>KO1</sup> allele through the germline was confirmed by Southern blot (upper panel) and a PCR genotyping assay (lower panel). PCR primers are indicated as arrows in A. Because the homozygous mutant embryos were being resorbed, there was much less yolk sac DNA in the -/- lanes.

Fig. 4 (panels A-G) illustrate the germ layer-specific derepression of Hh target genes in *ptc*<sup>-/-</sup> embryos. (A, B) Lateral views of E8.25 wild-type (A) and *ptc*<sup>-/-</sup> (B) embryos. The

headfolds are overgrown in the mutant (white arrows) and the heart is not properly formed (red arrows). (C) Lateral views of E8.75 *ptc*<sup>+/−</sup> (left) and *ptc*<sup>−/−</sup> (right) embryos stained with X-gal (28) (D, E, F, G) Transverse sections through E8.75 *ptc*<sup>+/−</sup> (D, F) and *ptc*<sup>−/−</sup> (E, G) embryos stained with X-gal (D, E) or hybridized with a digoxigenin labeled *Gli* probe (29) (F, G). Both *lacZ* and *Gli* were derepressed in the ectoderm and mesoderm but not in the endoderm (arrows). In A and B, anterior is to the left and dorsal is up. In C, anterior is up and dorsal is to the right. In D to G, dorsal is up.

Fig. 5 (panels A-L) illustrate ventralization of the neural tube in *ptc*<sup>−/−</sup> embryos. (A) Lateral view of E8.5 wild-type (left) and *ptc*<sup>−/−</sup> (right) embryos hybridized with a *HNF3b* probe. Expression is expanded dorsally in the mutant. (B, C) Transverse sections through the hindbrain of E8.5 wild-type (B) and *ptc*<sup>−/−</sup> (C) embryos hybridized with <sup>35</sup>S-labeled *Shh* probe (8). *Shh* is expressed in the floor plate (fp) and notochord (nc) of the wild-type embryo, and is greatly expanded in the *ptc* mutant. g = gut (D, E) Hematoxylin and eosin stained transverse sections through the hindbrain of wild-type (D) and *ptc*<sup>−/−</sup> (E) E8.5 embryos. Bottle-shaped cells with basal nuclei are indicated by arrows. (F, G) Transverse sections through E8.5 *ptc*<sup>+/−</sup> (F) and *ptc*<sup>−/−</sup> (G) embryos hybridized with *Pax6* probe show loss of expression from the *ptc* mutant. (H) Dorsal view of E8.25-E8.5 embryos hybridized with *Pax3* probe. Because of the kinking in the neural tube, the *ptc*<sup>−/−</sup> embryo is curled on itself. Weak *Pax3* expression is seen in the posterior dorsal neural tube of the *ptc*<sup>−/−</sup> embryo (bottom, arrow). (I, J) Transverse sections through E8.5 wild-type (I) and *ptc*<sup>−/−</sup> (J) embryos hybridized with *Pax3* probe. *Pax3* is expressed in the dorsal neural tube (nt) and dermamyotome (dm) in the wild-type, but is only present in a small dorsal domain of the mutant neural tube. s = somite (K, L) Lateral views of E9 wild type (K) and E8.5 *ptc*<sup>−/−</sup> (L) embryos hybridized with *erb-b3* probe. Staining is seen in migrating neural crest in the head and somites of wild type but not mutant embryos (red arrows). Weak staining in the head, heart and gut (black arrows) is background or non-neural crest related. (M) Lateral view of wild type (top) and *ptc*<sup>−/−</sup> (bottom) embryos hybridized with *Nkx2.1* probe. The body of the mutant is twisted. *Nkx2.1* expression is limited to the anterior, but is expanded dorsally in the mutant. (N) Lateral view of E8.5 *ptc*<sup>+/−</sup> (left) and *ptc*<sup>−/−</sup> (right) embryos hybridized with *hoxb1* probe. Loss of expression in rhombomere four is indicated by the asterisks. In all transverse sections, dorsal is up. In A, K, L and N, anterior is up and dorsal is to the right. In H and M, anterior is to the left.

Fig. 6 (panels A-F) depict keletal abnormalities and medulloblastomas in *ptc*<sup>+/−</sup> mice (A) Alcian blue and Alizarin red stained hindlimb from a *ptc*<sup>+/−</sup> mouse (30). The preaxial digit is duplicated (arrows). (B, C) Dorsal views of brains from wild-type (B) and *ptc*<sup>+/−</sup> (C) mice. Anterior is up. In the posterior wild-type brain, the colliculi (col) are present as distinct bumps between the cortex (cor) and cerebellum (ce). In the *ptc*<sup>+/−</sup> mouse, a massive medulloblastoma (mb, outlined in red) grew over the colliculi and normal

cerebellum, which can no longer be seen. The olfactory bulbs were removed. (D, E) Hematoxylin and eosin stained section through human (D) and mouse (E) medulloblastomas. The tumor cells are small with dark, carrot-shaped nuclei (arrows) and form nodules with no apparent orientation. (F) Synaptophysin immunoreactivity in a mouse medulloblastoma (26). Synaptophysin staining (brown) is seen in some processes (arrows). Nuclei are purple.

Fig. 7 (panels A-G) illustrate derepression of *ptc* and *Gli* expression in medulloblastomas from *ptc*<sup>+/-</sup> mice. (A to C) Semi-adjacent sections through a tumor in the cerebellum of a *ptc*<sup>+/-</sup> mouse hybridized with <sup>35</sup>S labeled probes to *ptc* (A), *Gli* (B) and *Shh* (C). *ptc* and *Gli* transcripts are abundant in the tumors (asterisks) compared to nearby cerebellar tissue (arrows). No *Shh* was detected in the tumor. (D) *ptc*<sup>+/-</sup> cerebellum (ce) and tumor (mb) stained with X-gal (28). Anterior is to the left. Derepression of *ptc* expression in the medulloblastoma is reflected in the high level of X-gal staining. (E) Surface staining in (arrows) regions of *ptc*<sup>+/-</sup> cerebellum contrast with absence of β-galactosidase activity in most folia (asterisk). (F) Sagittal section through cerebellum in E. X-gal staining nuclei (arrow) accumulated superficial to the molecular layer (ml), where stained nuclei are not normally seen. In unaffected regions of the cerebellum, X-gal staining was seen in scattered cells of the molecular layer (ml), strongly in the Purkinje cell layer (pcl) and weakly in the granule cell layer (gl). (G) *ptc* expression was examined in total RNA (15 mg) from wild-type (WT) and *ptc*<sup>+/-</sup> (+/-) cerebellums using a probe (M2-2) (6) that detects exons downstream of the *lacZ* and *neo* insertions. Actin mRNA was used as an RNA loading control. The *ptc*<sup>+/-</sup> mice had ~50% decrease in *ptc* transcripts.

#### Database References for Nucleotide and Amino Acid Sequences

The sequence for the *D. melanogaster patched* gene has the Genbank accession number M28418. The sequence for the mouse *patched* gene has the Genbank accession number U130589-V46155. The sequence for the human *patched* gene has the Genbank accession number U59464.

#### Detailed Description of the Invention

Vertebrate and invertebrate *patched* (*ptc*) gene compositions and methods for their isolation are provided. Of particular interest are mammalian *ptc* genes, such as the human and mouse homologs described in the appended examples. The *ptc* gene, in mammals, is a tumor suppressor and developmental regulator. Certain human cancers, e.g. basal cell carcinoma, transitional cell carcinoma of the bladder, meningiomas, medulloblastomas, etc., can be characterized by *ptc* loss-of-function, such as that resulting from oncogenic mutations at the *ptc* locus, or other loss-of-function mutations which decrease *ptc* activity in the cell. As

described below, we have observed somatic mutations in the *ptc* gene in a variety of sporadic cancers. For instance, the basal cell nevus syndrome (BCNS), an inherited disorder, is associated with germline mutations in *ptc*. Some patients with basal cell nevus syndrome (BCNS) have germ line mutations in *ptc*, and are at increased risk for developmental defects  
5 such as spina bifida and craniofacial abnormalities, basal cell carcinoma (BCC) of the skin, and brain tumors. Mutations to *ptc* genes are also observed to occur in sporadic BCCs, which generally have both copies of *ptc* inactivated.

The term “loss-of-function” is art recognized and, with respect to a *patched* gene or gene product refers to mutations in a *patched* gene which ultimately decrease or otherwise  
10 inhibit the ability of a cell to transduce *patched*-mediated signals, e.g., the cells may lose responsiveness to *hedgehog* induction. For example, a loss-of-function mutation to a *patched* gene may be a point mutation, deletion or insertion of sequences in the coding sequence, intron sequence or 5’ or 3’ flanking sequences of the gene so as to, for example, (i) alter (e.g., decrease) the level *patched* expression, (ii) alter exon-splicing patterns, (iii) alter the ability of  
15 the encoded *patched* protein to interact with extracellular or intracellular proteins (such as *hedgehog*), or (iv) alter (decrease) the stability of the encoded *patched* protein.

The term “aberrant modification” is art recognized and, with respect to a *patched* gene, refers to a non-wildtype mutation or other alteration to the gene, e.g., which results in full or partial loss-of-function of the *patched* protein or expression of the *patched* gene.

Such mutations affecting *ptc* activity have also been associated with other human  
20 cancers, including carcinomas, adenocarcinomas, sarcomas and the like. Decreased *ptc* activity is also associated with inherited developmental abnormalities, e.g. rib and craniofacial abnormalities, polydactyly, syndactyly and spina bifida.

The art-recognized term “predisposing mutation”, as it pertains to *patched* genes,  
25 refers to mutations to the *patched* gene which result in loss-of-function.

The term “genetic predisposition” is art recognized, and refers to a genotype of an animal which predisposes the animal to developing a certain pathological conditions with a frequency (probability) greater than the average for the overall population of that animal, taking into account, as appropriate, age, sex or other related physical or medical condition(s).

30 The *ptc* genes and fragments thereof, encoded protein, and anti-*ptc* antibodies are useful in the identification of individuals predisposed to development of a variety of cancers and developmental abnormalities, and in characterizing the phenotype of various tumors or other proliferative or degenerative disorders that are associated with this gene, e.g., for diagnostic and/or prognostic benefit. The characterization is useful for prenatal screening; and  
35 in determining the phenotype of a proliferative disorder, e.g. for determining a course of treatment of the patient. Tumors may be typed or staged as to the *ptc* status, e.g. by detection

of mutated sequences, antibody detection of abnormal protein products, and functional assays for altered *ptc* activity.

The terms “developmental disorder” and “developmental abnormality” are art recognized, and refer to aberrant development of a cell, tissue or organ, e.g., in size, symmetry or functional performance, which abnormality may or may not be untowardly manifest.

The term “proliferative disorder” is art recognized and refers to a disorder affecting an animal in a manner which is marked by aberrant, or otherwise unwanted, proliferation of a subset of cells of an animal. Cancers are proliferative disorders.

10 The encoded *ptc* protein is also useful in drug screening for compositions that mimic *ptc* activity or expression, including altered forms of *ptc* protein, particularly with respect to *ptc* function as a tumor suppressor in oncogenesis.

15 The human and mouse *ptc* gene sequences and isolated nucleic acid compositions are provided in the appended examples. In identifying the mouse and human *patched* genes, cross-hybridization of DNA and amplification primers were employed to move through the evolutionary tree from the known *Drosophila ptc* sequence, identifying a number of invertebrate homologs.

20 The human *patched* gene has been mapped to human chromosome band 9q22.3, and lies between the polymorphic markers D9S196 and D9S287 (a detailed map of human genome markers may be found in Dib *et al.* (1996) *Nature* 280:152- <http://www.genethon.fr>).

As will be understood by those skilled in the art, the method of the present invention can be carried out using any of a large number of assay techniques for detecting alterations in *ptc* genes and/or *ptc* protein function. For instance, individuals are screened by analyzing their DNA or RNA for the presence of a predisposing oncogenic or developmental mutation, as compared to a normal sequence. An exemplary "normal" sequence of *patched* is provided in SEQ ID NO:19 (human). Specific mutations of interest include any mutation that leads to oncogenesis or developmental abnormalities, including insertions, substitutions and deletions in the coding region sequence, in the introns (e.g., that affect splicing), in the transcriptional regulatory sequences (such as promoter or enhancer sequences) that affect the activity and expression of the protein.

In general, the subject method can be characterized as including a step of detecting, in a sample of cells from a patient, the presence or absence of *ptc* expression (at the protein or mRNA transcript level), mutations to the *ptc* gene (coding or non-coding sequence) and/or the functional activity of *ptc* in the sample of cells (such as induction of Gli or the like). Moreover, the subject method can be used to assess the phenotype of cells which are known

to be transformed, the phenotype results being useful in planning a particular therapeutic regimen.

To illustrate, nucleic acid samples are obtained from a patient having, or suspected as being at risk for developing, a tumor or developmental abnormality which may be associated with *ptc*. The nucleic acid is analyzed for the presence of a predisposing mutation in the *ptc* gene. The presence of a mutated *ptc* sequence that affects the level of expression of the gene, stability of the gene product, and/or signal transduction activity of *ptc* confers an increased susceptibility to a proliferative or developmental disorder. Thus, the level of expression of *ptc* can be used predictively to evaluate whether a sample of cells contains cells which are, or are predisposed towards becoming, transformed.

Diagnostic/prognostic screening of tissue/cell samples for tumors or developmental abnormalities may also be based on the functional or antigenic characteristics of the protein. Immunoassays designed to detect the normal or abnormal *ptc* protein may be used in screening. Where many diverse mutations lead to a particular disease phenotype, functional protein assays have proven to be effective screening tools. Such assays may be based on detecting changes in the transcriptional regulation mediated by *ptc*, or may directly detect *ptc* activities such as *hedgehog* binding, transporter activity or the like, or may involve antibody localization of *patched* in cells.

Inheritance of BCNS is autosomal dominant, although many cases are the result of new mutations. Diagnosis of BCNS is performed by protein, DNA sequence or hybridization analysis of any convenient sample from a patient, e.g. biopsy material, blood sample, scrapings from cheek, etc. A typical patient genotype will have a predisposing mutation on at least one chromosome. In tumors and at least sometimes developmentally affected tissues, loss of heterozygosity at the *ptc* locus leads to aberrant cell and tissue behavior. When the normal copy of *ptc* is lost, leaving only the reduced function mutant copy, abnormal cell growth and reduced cell layer adhesion is the result. Examples of specific *ptc* mutations in BCNS patients are a 9 bp insertion at nt 2445 of the coding sequence- and an 11 bp deletion of nt 2441 to 2452 of the coding sequence. These result in insertions or deletions in the region of the seventh transmembrane domain.

Prenatal diagnosis of BCNS may be performed, particularly where there is a family history of the disease, e.g. an affected parent or sibling. It is desirable, although not required, in such cases to determine the specific predisposing mutation present in affected family members. A sample of fetal DNA, such as an amniocentesis sample, fetal nucleated or white blood cells isolated from maternal blood, chorionic villus sample, etc. is analyzed for the presence of the predisposing mutation. Alternatively, a protein based assay, e.g. functional assay or immunoassay, is performed on fetal cells known to express *ptc*.

Sporadic tumors associated with loss of *ptc* function include a number of carcinomas and other transformed cells known to have deletions in the region of chromosome 9q22, e.g. basal cell carcinomas, transitional bladder cell carcinoma, meningiomas, medullomas, fibromas of the heart and ovary, and carcinomas of the lung, ovary, kidney and esophagus.

5 Characterization of sporadic tumors will generally require analysis of tumor cell DNA, conveniently with a biopsy sample. A wide range of mutations are found in sporadic cases, up to and including deletion of the entire long arm of chromosome 9. Oncogenic mutations may delete one or more exons, e.g. 8 and 9, may affect the amino acid sequence such as of the extracellular loops or transmembrane domains, may cause truncation of the protein by  
10 introducing a frameshift or stop codon, etc. Specific examples of oncogenic mutations include a C to T transition at nt 523 and deletions encompassing exon 9. C to T transitions are characteristic of ultraviolet mutagenesis, as expected with cases of skin cancer.

Biochemical studies may be performed to determine whether a candidate sequence variation in the *ptc* coding region or control regions is oncogenic. For example, a change in  
15 the promoter or enhancer sequence that downregulates expression of *patched* may result in predisposition to cancer. Expression levels of a candidate variant allele are compared to expression levels of the normal allele by various methods known in the art. Methods for determining promoter or enhancer strength include quantitation of the expressed mRNA or *ptc* protein; insertion of the variant control element into a vector with a reporter gene such as  
20  $\beta$ -galactosidase, chloramphenical acetyltransferase, etc. that provides for convenient quantitation- and the like. Nuclear run-off assays are another convenient means for measuring promoter/enhancer activity. The activity of the encoded *ptc* protein may be determined by comparison with the wild-type protein, e.g. by detection of transcriptional regulation of TGF or *Wnt* family genes, Gli genes, *ptc* itself, or reporter gene fusions  
25 involving transcriptional regulatory sequences of these target genes.

The term “*patched*-dependent gene”, or “a gene which is regulated in a *patched*-dependent manner”, refers to genes, such as Gli or *patched*, etc, whose level of expression is regulated at least in part by the presence of a *patched* protein in the cell, e.g., can be controlled by *patched*-dependent intracellular signals.

30 A human *patched* gene (SEQ ID NO:18) has a 4.5 kb open reading frame encoding a protein of 1447 amino acids. Including coding and noncoding sequences, it is about 89% identical at the nucleotide level to the mouse *patched* gene (SEQ ID NO:9). A mouse *patched* gene (SEQ ID NO:9) encodes a protein (SEQ ID NO:10) that has about 38% identical amino acids to *Drosophila ptc* (SEQ ID NO:6), over about 1,200 amino acids. The butterfly  
35 homolog (SEQ ID NO:4) is 1,300 amino acids long and overall has a 50% amino acid identity to fly *ptc* (SEQ ID NO:6). A 267 bp exon from the beetle *patched* gene encodes an

89 amino acid protein fragment, which was found to be 44% and 51% identical to the corresponding regions of fly and butterfly *ptc* respectively.

The DNA sequence encoding *ptc* may be cDNA, RNA, genomic DNA or synthetic, and includes fragments of the full-length coding sequence. The term "*patched* gene" shall be intended to mean the open reading frame encoding specific *ptc* polypeptides, as well as, as appropriate, adjacent intronic sequences and 5' and 3' non-coding nucleotide sequences involved in the regulation of expression, up to about 1 kb beyond the coding region, in either direction. The gene may be introduced into an appropriate vector for extrachromosomal maintenance or for integration into the host.

The term "cDNA" as used herein is intended to include all nucleic acids that share the arrangement of sequence elements found in native mature mRNA species, where sequence elements are exons, 3' and 5' non-coding regions. Normally mRNA species have contiguous exons, with the intervening introns deleted, to create a continuous open reading frame encoding *ptc*.

The genomic *ptc* sequence has a non-contiguous open reading frame, where introns interrupt the coding regions. A genomic sequence of interest comprises the nucleic acid present between the initiation codon and the stop codon, as defined in the listed sequences, including all of the introns that are normally present in a native chromosome. It may further include the 3' and 5' untranslated regions found in the mature mRNA. It may further include specific transcriptional and translational regulatory sequences, such as promoters, enhancers, etc., including about 1 kb of flanking genomic DNA at either the 5' or 3' end of the coding region. The genomic DNA may be isolated as a fragment of 50 kbp or smaller; and substantially free of flanking chromosomal sequence.

The nucleic acid compositions of the subject invention encode all or a part of the subject polypeptides. Fragments may be obtained of the DNA sequence by chemically synthesizing oligonucleotides in accordance with conventional methods, by restriction enzyme digestion, by PCR amplification, etc. For the most part, DNA fragments will be of at least 15 nt, usually at least 18 nt, more usually at least about 50 nt. Such small DNA fragments are useful as primers for PCR, hybridization screening, etc. Larger DNA fragments, i.e. greater than 100 nt are useful for production of the encoded polypeptide. For use in amplification reactions, such as PCR, a pair of primers will be used. The exact composition of the primer sequences is not critical to the invention, but for most applications the primers will hybridize to the subject sequence under stringent conditions, as known in the art. It is preferable to choose a pair of primers that will generate an amplification product of at least about 50 nt, preferably at least about 100 nt. Algorithms for the selection of primer sequences are generally known, and are available in commercial software packages.

Amplification primers hybridize to complementary strands of DNA, and will prime towards each other.

The *ptc* genes are isolated and obtained in substantial purity, generally as other than an intact mammalian chromosome. Usually, the DNA will be obtained substantially free of other nucleic acid sequences that do not include a *ptc* sequence or fragment thereof, generally being at least about 50%, usually at least about 90% pure and are typically "recombinant", i.e. flanked by one or more nucleotides with which it is not normally associated on a naturally occurring chromosome.

The DNA sequences are used in a variety of ways. They may be used as probes for identifying other patched genes. Mammalian homologs have substantial sequence similarity to the subject sequences, i.e. at least 75%, usually at least 90%, more usually at least 95% sequence identity with the nucleotide sequence of the subject DNA sequence. Sequence similarity is calculated based on a reference sequence, which may be a subset of a larger sequence, such as a conserved motif, coding region, flanking region, etc. A reference sequence will usually be at least about 18 nt long, more usually at least about 30 nt long, and may extend to the complete sequence that is being compared. Algorithms for sequence analysis are known in the art, such as BLAST, described in Altschul *et al.* (1990) J Mol Biol 215:403-10.

Nucleic acids having sequence similarity are detected by hybridization under low stringency conditions, for example, at 50 C and 10xSSC (0.9 M saline/0.09 M sodium citrate) and remain bound when subjected to washing at 55°C in 1xSSC. By using probes, particularly labeled probes of DNA sequences, one can isolate homologous or related genes. The source of homologous genes may be any mammalian species, e.g. primate species, particularly human- murines, such as rats and mice, canines, felines, bovines, ovines, equines, etc.

The DNA may also be used to identify expression of the gene in a biological specimen. The manner in which one probes cells for the presence of particular nucleotide sequences, as genomic DNA or RNA, is well-established in the literature and does not require elaboration here. Conveniently, a biological specimen is used as a source of mRNA. The mRNA may be amplified by RT-PCR, using reverse transcriptase to form a complementary DNA strand, followed by polymerase chain reaction amplification using primers specific for the subject DNA sequences. Alternatively, the mRNA sample is separated by gel electrophoresis, transferred to a suitable support, e.g., nitrocellulose and then probed with a fragment of the subject DNA as a probe. Other techniques may also find use. Detection of mRNA having the subject sequence is indicative of *patched* gene expression in the sample.

The subject nucleic acid sequences may be modified for a number of purposes, particularly where they will be used intracellularly, for example, by being joined to a nucleic acid cleaving agent, e.g. a chelated metal ion, such as iron or chromium for cleavage of the

gene; as an antisense sequence, or the like. Modifications may include replacing oxygen of the phosphate esters with sulfur or nitrogen, replacing the phosphate with phosphoramidate, etc.

A number of methods are available for analyzing genomic DNA sequences. Where large amounts of DNA are available, the genomic DNA is used directly. Alternatively, the region of interest is cloned into a suitable vector and grown in sufficient quantity for analysis, or amplified by conventional techniques, such as the polymerase chain reaction (PCR). The use of the polymerase chain reaction is described in Saiki, *et al.* (1985) Science 239:487, and a review of current techniques may be found in Sambrook, *et al.* Molecular Cloning: A Laboratory Manual, CSH Press 1989, pp.14.2-14.33.

A detectable label may be included in the amplification reaction. Suitable labels include fluorochromes, e.g. fluorescein isothiocyanate (FITC), rhodamine, Texas Red, phycoerythrin, allophycocyanin, 6-carboxyfluorescein (6-FAM), 2',7'-dimethoxy-4',5'-dichloro-6-carboxyfluorescein (JOE), 6-carboxy-Xrhodamine (ROX), 6-carboxy-2',4',7',4,7-hexachlorofluorescein (HEX), 5-carboxyfluorescein (5-FAM) or N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA), radioactive labels, e.g. **32P**, **35S**, **3H**; etc. The label may be a two stage system, where the amplified DNA is conjugated to biotin, haptens, etc. having a high affinity binding partner, e.g. avidin, specific antibodies, etc., where the binding partner is conjugated to a detectable label. The label may be conjugated to one or both of the primers. Alternatively, the pool of nucleotides used in the amplification is labeled, so as to incorporate the label into the amplification product.

The amplified or cloned fragment may be sequenced by dideoxy or other methods, and the sequence of bases compared to the normal *ptc* sequence. Hybridization with the variant sequence may also be used to determine its presence, by Southern blots, dot blots, etc. Single strand conformational polymorphism (SSCP) analysis, denaturing gradient gel electrophoresis (DGGE), and heteroduplex analysis in gel matrices are used to detect conformational changes created by DNA sequence variation as alterations in electrophoretic mobility. The hybridization pattern of a control and variant sequence to an array of oligonucleotide probes immobilized on a solid support, as described in WO 95/11995, may also be used as a means of detecting the presence of variant sequences. Alternatively, where a predisposing mutation creates or destroys a recognition site for a restriction endonuclease, the fragment is digested with that endonuclease, and the products size fractionated to determine whether the fragment was digested. Fractionation is performed by gel electrophoresis, particularly acrylamide or agarose gels.

In a merely illustrative embodiment, the method includes the steps of (i) collecting a sample of cells from a patient, (ii) isolating nucleic acid (e.g., genomic, mRNA or both) from the cells of the sample, (iii) contacting the nucleic acid sample with one or more primers

which specifically hybridize to a *ptc* gene under conditions such that hybridization and amplification of the *ptc* gene (if present) occurs, and (iv) detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample.

5 In yet another exemplary embodiment, aberrant methylation patterns of a *ptc* gene can be detected by digesting genomic DNA from a patient sample with one or more restriction endonucleases that are sensitive to methylation and for which recognition sites exist in the *ptc* gene (including in the flanking and intronic sequences). See, for example, Buiting et al., (1994) Human Mol Genet 3:893-895. Digested DNA is separated by gel electrophoresis, and  
10 hybridized with probes derived from, for example, genomic or cDNA sequences. The methylation status of the *ptc* gene can be determined by comparison of the restriction pattern generated from the sample DNA with that for a standard of known methylation.

In still another embodiment, a diagnostic assay is provided which detects the ability of a *ptc* gene product, e.g., recombinantly expressed from a gene isolated from a biopsied  
15 cell, to bind to other proteins, e.g., upstream (*hedgehog*) or downstream of *ptc*. For instance, it will be desirable to detect *ptc* mutants which bind with lower binding affinity for *hedgehog* proteins. Such mutants may arise, for example, from fine mutations, e.g., point mutants, which may be impractical to detect by the diagnostic DNA sequencing techniques or by the immunoassays described above. The present invention accordingly further contemplates  
20 diagnostic screening assays which generally comprise cloning one or more *ptc* genes from the sample cells, and expressing the cloned genes under conditions which permit detection of an interaction between that recombinant gene product and a *ptc*-binding protein, e.g., a *hedgehog* protein. As will be apparent from the description of the various drug screening assays set forth below, a wide variety of techniques can be used to determine the ability of a  
25 *ptc* protein to bind to other cellular components.

The subject nucleic acids can be used to generate transgenic animals or site specific gene modifications in cell lines. Transgenic animals may be made through homologous recombination, where the normal patched locus is altered. Alternatively, a nucleic acid construct is randomly integrated into the genome. Vectors for stable integration include  
30 plasmids, retroviruses and other animal viruses, YACS, and the like.

The modified cells or animals are useful in the study of *patched* function and regulation. For example, a series of small deletions and/or substitutions may be made in the *patched* gene to determine the role of different exons in oncogenesis, signal transduction, etc. Of particular interest are transgenic animal models for carcinomas of the skin, where  
35 expression of *ptc* is specifically reduced or absent in skin cells. An alternative approach to transgenic models for this disease are those where one of the mammalian *hedgehog* genes, e.g. *Shh*, *lh*, *Dhh*, are upregulated in skin cells, or in other cell types. For models of skin

abnormalities, one may use a skin-specific promoter to drive expression of the transgene, or other inducible promoter that can be regulated in the animal model. Such promoters include keratin gene promoters. Specific constructs of interest include anti-sense *ptc*, which will block *ptc* expression, expression of dominant negative *ptc* mutations, and over-expression of HH genes. A detectable marker, such as *lacZ* may be introduced into the *patched* locus, where upregulation of patched expression will result in an easily detected change in phenotype.

One may also provide for expression of the *patched* gene or variants thereof in cells or tissues where it is not normally expressed or at abnormal times of development. Thus, mouse models of spina bifida or abnormal motor neuron differentiation in the developing spinal cord are made available. In addition, by providing expression of *ptc* protein in cells in which it is otherwise not normally produced, one can induce changes in cell behavior, e.g. through *ptc* mediated transcription modulation.

DNA constructs for homologous recombination will comprise at least a portion of the *patched* or *hedgehog* gene with the desired genetic modification, and will include regions of homology to the target locus. DNA constructs for random integration need not include regions of homology to mediate recombination. Conveniently, markers for positive and negative selection are included. Methods for generating cells having targeted gene modifications through homologous recombination are known in the art. For various techniques for transfecting mammalian cells, see Keown *et al.* (1990) Methods in Enzymology 185:527-537.

For embryonic stem (ES) cells, an ES cell line may be employed, or ES cells may be obtained freshly from a host, e.g. mouse, rat, guinea pig, etc. Such cells are grown on an appropriate fibroblast-feeder layer or grown in the presence of leukemia inhibiting factor (LIF). When ES cells have been transformed, they may be used to produce transgenic animals. After transformation, the cells are plated onto a feeder layer in an appropriate medium. Cells containing the construct may be detected by employing a selective medium. After sufficient time for colonies to grow, they are picked and analyzed for the occurrence of homologous recombination or integration of the construct. Those colonies that are positive may then be used for embryo manipulation and blastocyst injection. Blastocysts are obtained from 4 to 6 week old superovulated females. The ES cells are trypsinized, and the modified cells are injected into the blastocoel of the blastocyst. After injection, the blastocysts are returned to each uterine horn of pseudopregnant females. Females are then allowed to go to term and the resulting litters screened for mutant cells having the construct. By providing for a different phenotype of the blastocyst and the ES cells, chimeric progeny can be readily detected.

The chimeric animals are screened for the presence of the modified gene and males and females having the modification are mated to produce homozygous progeny. If the gene alterations cause lethality at some point in development, tissues or organs can be maintained as allogeneic or congenic grafts or transplants, or in *in vitro* culture. The transgenic animals may be any non-human mammal, such as laboratory animals, domestic animals, etc. The transgenic animals may be used in functional studies, drug screening, etc., e.g. to determine the effect of a candidate drug on basal cell carcinomas.

The subject gene may be employed for producing all or portions of the patched protein. For expression, an expression cassette may be employed, providing for a transcriptional and translational initiation region, which may be inducible or constitutive, the coding region under the transcriptional control of the transcriptional initiation region, and a transcriptional and translational termination region. Various transcriptional initiation regions may be employed which are functional in the expression host.

Specific *ptc* peptides of interest include the extracellular domains, particularly in the human mature protein, aa 120 to 437, and aa 770 to 1027. These peptides may be used as immunogens to raise antibodies that recognize the protein in an intact cell membrane. The cytoplasmic domains, as shown in Figure 2, (the amino terminus and carboxy terminus) are of interest in binding assays to detect ligands involved in signaling mediated by *ptc*.

The peptide may be expressed in prokaryotes or eukaryotes in accordance with conventional ways, depending upon the purpose for expression. For large scale production of the protein, a unicellular organism or cells of a higher organism, e.g. eukaryotes such as vertebrates, particularly mammals, may be used as the expression host, such as *E. coli*, *B. subthis*, *S. cerevisiae*, and the like. In many situations, it may be desirable to express the *patched* gene in a mammalian host, whereby the *patched* gene will be glycosylated, and transported to the cellular membrane for various studies.

With the availability of the protein in large amounts by employing an expression host, the protein may be isolated and purified in accordance with conventional ways. A lysate may be prepared of the expression host and the lysate purified using HPLC, exclusion chromatography, gel electrophoresis, affinity chromatography, or other purification technique. The purified protein will generally be at least about 80% pure, preferably at least about 90% pure, and may be up to and including 100% pure. By pure is intended free of other proteins, as well as cellular debris.

The polypeptide is used for the production of antibodies, where short fragments provide for antibodies specific for the particular polypeptide, whereas larger fragments or the entire gene allow for the production of antibodies over the surface of the polypeptide or protein. Antibodies may be raised to the normal or mutated forms of *ptc*- The extracellular domains of the protein are of interest as epitopes, particular antibodies that recognize

common changes found in abnormal, oncogenic *ptc*, which compromise the protein activity. Antibodies may be raised to isolated peptides corresponding to these domains, or to the native protein, e.g. by immunization with cells expressing *ptc*, immunization with liposomes having *ptc* inserted in the membrane, etc. Antibodies that recognize the extracellular domains of *ptc* are useful in diagnosis, typing and staging of human carcinomas.

Antibodies are prepared in accordance with conventional ways, where the expressed polypeptide or protein may be used as an immunogen, by itself or conjugated to known immunogenic carriers, e.g. KLH, pre-S HBsAg, other viral or eukaryotic proteins, or the like. Various adjuvants may be employed, with a series of injections, as appropriate, For monoclonal antibodies, after one or more booster injections, the spleen may be isolated, the splenocytes immortalized, and then screened for high affinity antibody binding. The immortalized cells, e.g. hybridomas, producing the desired antibodies may then be expanded. For further description, see Monoclonal Antibodies- A Laboratory Manual, Harlow and Lane eds., Cold Spring Harbor Laboratories, Cold Spring Harbor, New York, 1988. If desired, the mRNA encoding the heavy and light chains may be isolated and mutagenized by cloning in *E. coli*, and the heavy and light chains may be mixed to further enhance the affinity of the antibody.

The antibodies find particular use in diagnostic assays for developmental abnormalities, basal cell carcinomas and other tumors associated with mutations in *ptc*. Staging, detection and typing of tumors may utilize a quantitative immunoassay for the presence or absence of normal *ptc*. Alternatively, the presence of mutated forms of *ptc* may be determined. A reduction in normal *ptc* and/or presence of abnormal *ptc* is indicative that the tumor is *ptc*-associated.

A sample is taken from a patient suspected of having a *ptc*-associated tumor, developmental abnormality or BCNS. Samples, as used herein, include biological fluids such as blood, cerebrospinal fluid, tears, saliva, lymph, dialysis fluid and the like- organ or tissue culture derived fluids, and fluids extracted from physiological tissues. Also included in the term are derivatives and fractions of such fluids. Biopsy samples are of particular interest, e.g. skin lesions, organ tissue fragments, etc. Where metastasis is suspected, blood samples may be preferred. The number of cells in a sample will generally be at least about  $10^3$ , usually at least  $10^4$  more usually at least about  $10^5$ . The cells may be dissociated, in the case of solid tissues, or tissue sections may be analyzed. Alternatively a lysate of the cells may be prepared.

Diagnosis may be performed by a number of methods. The different methods all determine the absence or presence of normal or abnormal *ptc* in patient cells suspected of having a mutation in *ptc*. For example, detection may utilize staining of intact cells or histological sections, performed in accordance with conventional methods. The antibodies of

interest are added to the cell sample, and incubated for a period of time sufficient to allow binding to the epitope, usually at least about 10 minutes. The antibody may be labeled with radioisotopes, enzymes, fluorescers, chemiluminescers, or other labels for direct detection. Alternatively, a second stage antibody or reagent is used to amplify the signal. Such reagents are well-known in the art. For example, the primary antibody may be conjugated to biotin, with horseradish peroxidase-conjugated avidin added as a second stage reagent. Final detection uses a substrate that undergoes a color change in the presence of the peroxidase. The absence or presence of antibody binding may be determined by various methods, including flow cytometry of dissociated cells, microscopy, radiography, scintillation counting, etc.

An alternative method for diagnosis depends on the *in vitro* detection of binding between antibodies and *ptc* in a lysate. Measuring the concentration of *ptc* binding in a sample or fraction thereof may be accomplished by a variety of specific assays. A conventional sandwich type assay may be used. For example, a sandwich assay may first attach *ptc*-specific antibodies to an insoluble surface or support. The particular manner of binding is not crucial so long as it is compatible with the reagents and overall methods of the invention. They may be bound to the plates covalently or non-covalently, preferably non-covalently.

The insoluble supports may be any compositions to which polypeptides can be bound, which is readily separated from soluble material, and which is otherwise compatible with the overall method. The surface of such supports may be solid or porous and of any convenient shape. Examples of suitable insoluble supports to which the receptor is bound include beads, e.g. magnetic beads, membranes and microtiter plates. These are typically made of glass, plastic (e.g. polystyrene), polysaccharides, nylon or nitrocellulose. Microtiter plates are especially convenient because a large number of assays can be carried out simultaneously, using small amounts of reagents and samples.

Patient sample lysates are then added to separately assayable supports (for example, separate wells of a microtiter plate) containing antibodies. Preferably, a series of standards, containing known concentrations of normal and/or abnormal *ptc* is assayed in parallel with the samples or aliquots thereof to serve as controls. Preferably, each sample and standard will be added to multiple wells so that mean values can be obtained for each. The incubation time should be sufficient for binding, generally, from about 0.1 to 3 hr is sufficient. After incubation, the insoluble support is generally washed of non-bound components. Generally, a dilute non-ionic detergent medium at an appropriate pH, generally 7-8, is used as a wash medium. From one to six washes may be employed, with sufficient volume to thoroughly wash nonspecifically bound proteins present in the sample.

After washing, a solution containing a second antibody is applied. The antibody will bind *ptc* with sufficient specificity such that it can be distinguished from other components present. The second antibodies may be labeled to facilitate direct, or indirect quantification of binding. Examples of labels that permit direct measurement of second receptor binding include radiolabels, such as  $^3\text{H}$  or  $^{125}\text{I}$ , fluorescers, dyes, beads, chemiluminescers, colloidal particles, and the like. Examples of labels which permit indirect measurement of binding include enzymes where the substrate may provide for a colored or fluorescent product. In a preferred embodiment, the antibodies are labeled with a covalently bound enzyme capable of providing a detectable product signal after addition of suitable substrate. Examples of suitable enzymes for use in conjugates include horseradish peroxidase, alkaline phosphatase, malate dehydrogenase and the like. Where not commercially available, such antibody-enzyme conjugates are readily produced by techniques known to those skilled in the art. The incubation time should be sufficient for the labeled ligand to bind available molecules. Generally, from about 0.1 to 3 hr is sufficient, usually 1 hr sufficing.

After the second binding step, the insoluble support is again washed free of non-specifically bound material. The signal produced by the bound conjugate is detected by conventional means. Where an enzyme conjugate is used, an appropriate enzyme substrate is provided so a detectable product is formed.

Other immunoassays are known in the art and may find use as diagnostics. Ouchterlony plates provide a simple determination of antibody binding. Western blots may be performed on protein gels or protein spots on filters, using a detection system specific for *ptc* as desired, conveniently using a labeling method as described for the sandwich assay.

Other diagnostic assays of interest are based on the functional properties of *ptc* protein itself. Such assays are particularly useful where a large number of different sequence changes lead to a common phenotype, i.e., loss of protein function leading to oncogenesis or developmental abnormality. For example, a functional assay may be based on the transcriptional changes mediated by *hedgehog* and *patched* gene products. Addition of soluble Hh to embryonic stem cells causes induction of transcription in target genes. The presence of functional *ptc* can be determined by its ability to antagonize Hh activity. Other functional assays may detect the transport of specific molecules mediated by *ptc*, in an intact cell or membrane fragment. Conveniently, a labeled substrate is used, where the transport in or out of the cell can be quantitated by radiography, microscopy, flow cytometry, spectrophotometry, etc. Other assays may detect conformational changes, or changes in the subcellular localization of *patched* protein.

By providing for the production of large amounts of patched protein, one can identify ligands or substrates that bind to, modulate or mimic the action of *patched*. A common feature in basal cell carcinoma is the loss of adhesion between epidermal and dermal layers,

indicating a role for *ptc* in maintaining appropriate cell adhesion. Areas of investigation include the development of cancer treatments, wound healing, adverse effects of aging, metastasis, etc.

Drug screening identifies agents that provide a replacement for *ptc* function in abnormal cells. The role of *ptc* as a tumor suppressor indicates that agents which mimic its function, in terms of transmembrane transport of molecules, transcriptional down-regulation, etc., will inhibit the process of oncogenesis. These agents may also promote appropriate cell adhesion in wound healing and aging, to reverse the loss of adhesion observed in metastasis, etc. Conversely, agents that reverse *ptc* function may stimulate controlled growth and healing. Of particular interest are screening assays for agents that have a low toxicity for human cells. A wide variety of assays may be used for this purpose, including labeled *in vitro* protein-protein binding assays, electrophoretic mobility shift assays, immunoassays for protein binding, and the like. The purified protein may also be used for determination of three-dimensional crystal structure, which can be used for modeling intermolecular interactions, transporter function, etc.

The term "agent" as used herein describes any molecule, e.g. protein or pharmaceutical, with the capability of altering or mimicking the physiological function of *patched*. Generally a plurality of assay mixtures are run in parallel with different agent concentrations to obtain a differential response to the various concentrations. Typically, one of these concentrations serves as a negative control, i.e. at zero concentration or below the level of detection.

Candidate agents encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 50 and less than about 2,500 daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or a combinations thereof.

Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides and oligopeptides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are

readily modified through conventional chemical, physical and biochemical means, and may be used to produce combinatorial libraries. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification, etc. to produce structural analogs.

5 Where the screening assay is a binding assay, one or more of the molecules may be joined to a label, where the label can directly or indirectly provide a detectable signal. Various labels include radioisotopes, fluorescers, chemilumescers, enzymes, specific binding molecules, particles, e.g. magnetic particles, and the like. Specific binding molecules include pairs, such as biotin and streptavidin, digoxin and antidigoxin etc. For the specific  
10 binding members, the complementary member would normally be labeled with a molecule that provides for detection, in accordance with known procedures.

A variety of other reagents may be included in the screening assay. These include reagents like salts, neutral proteins, e.g. albumin, detergents, etc. that are used to facilitate optimal protein-protein binding and/or reduce nonspecific or background interactions.  
15 Reagents that improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc. may be used. The mixture of components are added in any order that provides for the requisite binding. Incubations are performed at any suitable temperature, typically between 4° and 40° C. Incubation periods are selected for optimum activity, but may also be optimized to facilitate rapid high-throughput screening. Typically  
20 between 0.1 and 1 hours will be sufficient.

Other assays of interest detect agents that mimic patched function, such as repression of target gene transcription, transport of patched substrate compounds, etc. For example, an expression construct comprising a *patched* gene may be introduced into a cell line under conditions that allow expression. The level of patched activity is determined by a functional  
25 assay, as previously described. In one screening assay, candidate agents are added in combination with a Hh protein, and the ability to overcome Hh antagonism of *ptc* is detected. In another assay, the ability of candidate agents to enhance *ptc* function is determined. Alternatively, candidate agents are added to a cell that lacks functional *ptc*, and screened for the ability to reproduce *ptc* in a functional assay.

30 In one embodiment, the drug screening assay is a cell-based assay which detects the ability of a compound to alter *patched*-dependent gene transcription. By selecting transcriptional regulatory sequences from genes whose expression is regulated by *patched* signal transduction, e.g. from *patched*, *GLI*, *hedgehog* or PTHrP genes, e.g., regulatory sequences that are responsible for the up- or down regulation of these genes in response to  
35 *patched* signalling, and operatively linking such promoters to a reporter gene, one can derive a transcription based assay which is sensitive to the ability of a specific test compound to modify *patched* signalling pathways. Expression of the reporter gene, thus, provides a

valuable screening tool for the development of compounds that act as agonists or antagonists of *patched*.

Reporter gene based assays of this invention measure the end stage of the above described cascade of events, e.g., transcriptional modulation. Accordingly, in practicing one embodiment of the assay, a reporter gene construct is inserted into the reagent cell in order to generate a detection signal dependent on *ptc* signaling. To identify potential regulatory elements responsive to *ptc* signaling present in the transcriptional regulatory sequence of a target gene, nested deletions of genomic clones of the target gene can be constructed using standard techniques. See, for example, Current Protocols in Molecular Biology, Ausubel, F.M. et al. (eds.) Greene Publishing Associates, (1989); U.S. Patent 5,266,488; Sato et al. (1995) *J Biol Chem* 270:10314-10322; and Kube et al. (1995) *Cytokine* 7:1-7. A nested set of DNA fragments from the gene's 5'-flanking region are placed upstream of a reporter gene, such as the luciferase gene, and assayed for their ability to direct reporter gene expression in *patched* expressing cells. Host cells transiently transfected with reporter gene constructs can be scored for the induction of expression of the reporter gene in the presence and absence of *hedgehog* to determine regulatory sequences which are responsive to *patched*-dependent signalling.

In practicing one embodiment of the assay, a reporter gene construct is inserted into the reagent cell in order to generate a detection signal dependent on second messengers generated by induction with *hedgehog* protein. Typically, the reporter gene construct will include a reporter gene in operative linkage with one or more transcriptional regulatory elements responsive to the *hedgehog* activity, with the level of expression of the reporter gene providing the *hedgehog*-dependent detection signal. The amount of transcription from the reporter gene may be measured using any method known to those of skill in the art to be suitable. For example, mRNA expression from the reporter gene may be detected using RNase protection or RNA-based PCR, or the protein product of the reporter gene may be identified by a characteristic stain or an intrinsic activity. The amount of expression from the reporter gene is then compared to the amount of expression in either the same cell in the absence of the test compound (or *hedgehog*) or it may be compared with the amount of transcription in a substantially identical cell that lacks the target receptor protein. Any statistically or otherwise significant difference in the amount of transcription indicates that the test compound has in some manner altered the signal transduction of the *patched* protein, e.g., the test compound is a potential *ptc* therapeutic.

As described in further detail below, in preferred embodiments the gene product of the reporter is detected by an intrinsic activity associated with that product. For instance, the reporter gene may encode a gene product that, by enzymatic activity, gives rise to a detection signal based on color, fluorescence, or luminescence. In other preferred embodiments, the

reporter or marker gene provides a selective growth advantage, e.g., the reporter gene may enhance cell viability, relieve a cell nutritional requirement, and/or provide resistance to a drug.

Preferred reporter genes are those that are readily detectable. The reporter gene may also be included in the construct in the form of a fusion gene with a gene that includes desired transcriptional regulatory sequences or exhibits other desirable properties. Examples of reporter genes include, but are not limited to CAT (chloramphenicol acetyl transferase) (Alton and Vapnek (1979), Nature 282: 864-869) luciferase, and other enzyme detection systems, such as beta-galactosidase; firefly luciferase (deWet et al. (1987), Mol. Cell. Biol. 7:725-737); bacterial luciferase (Engebrecht and Silverman (1984), PNAS 1: 4154-4158; Baldwin et al. (1984), Biochemistry 23: 3663-3667); alkaline phosphatase (Toh et al. (1989) Eur. J. Biochem. 182: 231-238, Hall et al. (1983) J. Mol. Appl. Gen. 2: 101), human placental secreted alkaline phosphatase (Cullen and Malim (1992) Methods in Enzymol. 216:362-368).

Transcriptional control elements which may be included in a reporter gene construct include, but are not limited to, promoters, enhancers, and repressor and activator binding sites. Suitable transcriptional regulatory elements may be derived from the transcriptional regulatory regions of genes whose expression is induced after modulation of a *patched* signal transduction pathway. The characteristics of preferred genes from which the transcriptional control elements are derived include, but are not limited to, low or undetectable expression in quiescent cells, rapid induction at the transcriptional level within minutes of extracellular simulation, induction that is transient and independent of new protein synthesis, subsequent shut-off of transcription requires new protein synthesis, and mRNAs transcribed from these genes have a short half-life. It is not necessary for all of these properties to be present.

The compounds having the desired pharmacological activity may be administered in a physiologically acceptable carrier to a host for treatment of cancer or developmental abnormalities attributable to a defect in *patched* function. The compounds may also be used to enhance *patched* function in wound healing, aging, etc. The inhibitory agents may be administered in a variety of ways, orally, topically, parenterally e.g. subcutaneously, intraperitoneally, by viral infection, intravascularly, etc. Topical treatments are of particular interest. Depending upon the manner of introduction, the compounds may be formulated in a variety of ways. The concentration of therapeutically active compound in the formulation may vary from about 0.1-100 wt.%.

The pharmaceutical compositions can be prepared in various forms, such as granules, tablets, pills, suppositories, capsules, suspensions, salves, lotions and the like. Pharmaceutical grade organic or inorganic carriers and/or diluents suitable for oral and topical use can be used to make up compositions containing the therapeutically-active compounds. Diluents known to the art include aqueous media, vegetable and animal oils and fats. Stabilizing

agents, wetting and emulsifying agents, salts for varying the osmotic pressure or buffers for securing an adequate pH value, and skin penetration enhancers can be used as auxiliary agents.

The gene or fragments thereof may be used as probes for identifying the 5' non-coding region comprising the transcriptional initiation region, particularly the enhancer regulating the transcription of *patched*. By probing a genomic library, particularly with a probe comprising the 5' coding region, one can obtain fragments comprising the 5' non-coding region. If necessary, one may walk the fragment to obtain further 5' sequence to ensure that one has at least a functional portion of the enhancer. It is found that the enhancer is proximal to the 5' coding region, a portion being in the transcribed sequence and downstream from the promoter sequences. The transcriptional initiation region may be used for many purposes, studying embryonic development, providing for regulated expression of *patched* protein or other protein of interest during embryonic development or thereafter, and in gene therapy.

The gene may also be used for gene therapy. Vectors useful for introduction of the gene include plasmids and viral vectors. Of particular interest are retroviral-based vectors, e.g. moloney murine leukemia virus and modified human immunodeficiency virus-adenovirus vectors, etc. Gene therapy may be used to treat skin lesions, an affected fetus, etc., by transfection of the normal gene into embryonic stem cells or into other fetal cells. A wide variety of viral vectors can be employed for transfection and stable integration of the gene into the genome of the cells. Alternatively, micro-injection may be employed, fusion, or the like for introduction of genes into a suitable host cell. See, for example, Dhawan *et al.* (1991) Science 254:1509-1512 and Smith *et al.* (1990) Molecular and Cellular Biology 3268-3271.

The following examples are offered by illustration not by way of limitation.

## EXPERIMENTAL

### 25 Methods and Materials

*PCR on Mosquito (Anopheles gambiae) Genomic DNA.* PCR primers were based on amino acid stretches of fly *ptc* that were not likely to diverge over evolutionary time and were of low degeneracy. Two such primers (P2R1 (SEQ ID NO:14)-GGACGAATTCAARGTNCAYCARYTNTGG, P4R1: (SEQ ID NO:15)  
30 GGACGAATTCCYTCCCARAARCANTC, (the underlined sequences are Eco RI linkers) amplified an appropriately sized band from mosquito genomic DNA using the PCR. The program conditions were as follows:

94 C 4 min.; 72 C Add Taq;

[49 C 30 sec.; 72 C 90 sec.; 94 C 15 sec] 3 times

35 [94 C 15 sec.; 50 C 30 sec.; 72 C 90 sec] 35 times

72 C 10 min; 4 C hold

This band was subcloned into the EcoRV site of pBluescript II and sequenced using the USB Sequence kit.

5        *Screen of a Butterfly cDNA Library with Mosquito PCR Product.* Using the mosquito PCR product (SEQ ID NO:7) as a probe, a 3 day embryonic *Precis coenia* gt10 cDNA library (generously provided by Sean Carroll) was screened. Filters were hybridized at 65° C overnight in a solution containing 5xSSC, 10% dextran sulfate, 5x Denhardt's, 200 µg/ml sonicated salmon sperm DNA, and 0.5% SDS. Filters were washed in 0.1X SSC, 0.1% SDS  
10 at room temperature several times to remove nonspecific hybridization. Of the 100,000 plaques initially screened, 2 overlapping clones, L1 and L2, were isolated, which corresponded to the N terminus of butterfly *ptc*. Using L2 as a probe, the library filters were rescreened and 3 additional clones (L5, L7, L8) were isolated which encompassed the remainder of the *ptc* coding sequence. The full length sequence of butterfly *ptc* (SEQ ID  
15 NO:3) was determined by ABI automated sequencing.

*Screen of a Tribolium (beetle) Genomic Library with Mosquito PCR Product and 900 bp Fragment from the Butterfly Clone.* A *gem11* genomic library from *Tribolium castaneum* (gift of Rob Dennell) was probed with a mixture of the mosquito PCR (SEQ ID NO:7) product and BstXI/EcoRI fragment of L2. Filters were hybridized at 55 C overnight  
20 and washed as above. Of the 75,000 plaques screened, 14 clones were identified and the SacI fragment of T8 (SEQ ID NO:1), which crosshybridized with the mosquito and butterfly probes, was subcloned into pBluescript.

*PCR on Mouse cDNA Using Degenerate Primers Derived from Regions Conserved in the Four Insect Homologues.* Two degenerate PCR primers (P4REV- (SEQ ID NO:16)  
25 GGACGAATTCYTNGANTGYTTYTGGA- P22- (SEQ ID NO:17) CATACCAGCCAAG  
CTTGTCIGGCCARTGCAT) were designed based on a comparison of *ptc* amino acid sequences from fly (*Drosophila melanogaster*) (SEQ ID NO:6), mosquito (*Anopheles gambiae*) (SEQ ID NO:8), butterfly (*Precis coenia*) (SEQ ID NO:4), and beetle (*Tribolium castaneum*) (SEQ ID NO:2). I represents inosine, which can form base pairs with all four  
30 nucleotides. P22 was used to reverse transcribe RNA from 12.5 dpc mouse limb bud (gift from David Kingsley) for 90 min at 37 C. PCR using P4REV (SEQ ID NO:17) and P22 (SEQ ID NO:18) was then performed on 1 µl of the resultant cDNA under the following conditions:

94 C 4 min.; 72 C Add Taq;

35 [94 C 15 sec.- 50 C 30 sec.- 72 C 90 sec.] 35 times

72 C 10 min.-, 4 C hold

PCR products of the expected size were subcloned into the TA vector (Invitrogen) and sequenced with the Sequenase Version 2.0 DNA Sequencing Kit (U. S. B.).

Using the cloned mouse PCR fragment as a probe, 300,000 plaques of a mouse 8.5  
 5 dpc *gt10* cDNA library (a gift from Brigid Hogan) were screened at 65° C as above and  
 washed in 2x SSC, 0.1% SDS at room temperature. 7 clones were isolated, and three (M2,  
 M4, and M8) were subcloned into pBluescript II. 200,000 plaques of this library were  
 rescreened using first, a 1.1 kb *EcoRI* fragment from M2 to identify 6 clones (M9-M16) and  
 secondly a mixed probe containing the most N terminal (*XhoI* fragment from M2) and most C  
 10 terminal sequences (*BamHI/BglIII* fragment from M9) to isolate 5 clones (M17-M21). M9,  
 M10, M14, and M17-21 were subcloned into the *EcoRI* site of pBluescript II (Stratagene).

*RNA Blots and in situ Hybridizations in Whole and Sectioned Mouse Embryos:*

*Northerns.* A mouse embryonic Northern blot and an adult multiple tissue Northern  
 blot (obtained from Clontech) were probed with a 900 bp *EcoRI* fragment from an N terminal  
 15 coding region of mouse *ptc*. Hybridization was performed at 65° C in 5x SSPE, 10x  
 Denhardt's, 100 µg/ml sonicated salmon sperm DNA, and 2% SDS. After several short room  
 temperature washes in 2x SSC, 0.05% SDS, the blots were washed at high stringency in 0.1  
 X SSC, 0.1% SDS at 50° C.

*In situ hybridization of sections:* 7.75, 8.5, 11.5, and 13.5 dpc mouse embryos were  
 20 dissected in PBS and frozen in Tissue-Tek medium at -80° C. 12-16 µm frozen sections were  
 cut, collected onto VectaBond (Vector Laboratories) coated slides, and dried for 30-60  
 minutes at room temperature. After a 10 minute fixation in 4% paraformaldehyde in PBS, the  
 slides were washed 3 times for 3 minutes in PBS, acetylated for 10 minutes in 0.25% acetic  
 anhydride in triethanolamine, and washed three more times for 5 minutes in PBS.  
 25 Prehybridization (50% formamide, 5X SSC, 250 µg/ml yeast tRNA, 500 µg/ml sonicated  
 salmon sperm DNA, and 5x Denhardt's) was carried out for 6 hours at room temperature in  
 50% formamide/5x SSC humidified chambers. The probe, which consisted of 1 kb from the  
 N-terminus of *ptc*, was added at a concentration of 200-1000 ng/ml into the same solution  
 used for prehybridization, and then denatured for five minutes at 80° C. Approximately 75  
 30 µl of probe were added to each slide and covered with Parafilm. The slides were incubated  
 overnight at 65° C in the same humidified chamber used previously. The following day, the  
 probe was washed successively in 5X SSC (5 minutes, 65° C), 0.2X SSC (1 hour, 65° C), and  
 0.2X SSC (10 minutes, room temperature). After five minutes in buffer B1 (0.1M maleic acid,  
 0.15 M NaCl, pH 7.5), the slides were blocked for 1 hour at room temperature in 1% blocking  
 35 reagent (Boehringer-Mannheim) in buffer B1, and then incubated for 4 hours in buffer B1  
 containing the DIG-AP conjugated antibody (Boehringer-Mannheim) at a 1:5000 dilution.  
 Excess antibody was removed during two 15 minute washes in buffer B1, followed by five

minutes in buffer B3 (100 mM Tris, 100mM NaCl, 5mM MgCl<sub>2</sub>, pH 9.5). The antibody was detected by adding an alkaline phosphatase substrate (350 µl 75 mg/ml X-phosphate in DMF, 450 µl 50 mg/ml NBT in 70% DMF in 100 mls of buffer B3) and allowing the reaction to proceed overnight in the dark. After a brief rinse in 10 mM Tris, 1mM EDTA, pH 8.0, the slides were mounted with Aquamount (Lerner Laboratories).

*Drosophila 5-transcriptional initiation region -gal constructs.* A series of constructs were designed that link different regions of the *ptc* promoter from *Drosophila* to a LacZ reporter gene in order to study the cis regulation of the *ptc* expression pattern. See Fig. 1. A 10.8kb BamHI/BspMI fragment comprising the 5'-non-coding region of the mRNA at its 3'-terminus was obtained and truncated by restriction enzyme digestion as shown in Fig. 1. These expression cassettes were introduced into *Drosophila* lines using a P-element vector (Thummel *et al.* (1988) *Gene* 74:445-456), which were injected into embryos, providing flies which could be grown to produce embryos. (See Spradling and Rubin (1982) *Science* 218:341-347 for a description of the procedure.) The vector used a pUC8 background into which was introduced the white gene to provide for yellow eyes, portions of the P-element for integration, and the constructs were inserted into a polylinker upstream from the LacZ gene. The resulting embryos, larvae, and adults were stained using antibodies to LacZ protein conjugated to HRP and the samples developed with OPD dye to identify the expression of the LacZ gene. The staining pattern in embryos is described in Fig. 1, indicating whether there was staining during the early and late development of the embryo.

*Isolation of a Mouse ptc Gene.* Homologues of fly *ptc* (SEQ ID NO:6) were isolated from three insects: mosquito, butterfly and beetle, using either PCR or low stringency library screens. PCR primers to six amino acid stretches of *ptc* of low mutatability and degeneracy were designed. One primer pair, P2 and P4, amplified an homologous fragment of *ptc* from mosquito genomic DNA that corresponded to the first hydrophilic loop of the protein. The 345bp PCR product (SEQ ID NO:7) was subcloned and sequenced and when aligned to fly *ptc*, showed 67% amino acid identity.

The cloned mosquito fragment was used to screen a butterfly gt 10 cDNA library. Of 100,000 plaques screened, five overlapping clones were isolated and used to obtain the full length coding sequence. The butterfly *ptc* homologue (SEQ ID NO:4) is 1,311 amino acids long and overall has 50% amino acid identity (72% similarity) to fly *ptc*. With the exception of a divergent C-terminus, this homology is evenly spread across the coding sequence. The mosquito PCR clone (SEQ ID NO:7) and a corresponding fragment of butterfly cDNA were used to screen a beetle *gemll* genomic library. Of the plaques screened, 14 clones were identified. A fragment of one clone (T8), which hybridized with the original probes, was subcloned and sequenced. This 3kb piece contains an 89 amino acid exon (SEQ ID NO:2)

which is 44% and 51% identical to the corresponding regions of fly and butterfly *ptc* respectively.

Using an alignment of the four insect homologues in the first hydrophilic loop of the *ptc*, two PCR primers were designed to a five and six amino acid stretch which were identical and of low degeneracy. These primers were used to isolate the mouse homologue using RT-PCR on embryonic limb bud RNA. An appropriately sized band was amplified and upon cloning and sequencing, it was found to encode a protein 65% identical to fly *ptc*. Using the cloned PCR product and subsequently, fragments of mouse *ptc* cDNA, a mouse embryonic cDNA library was screened. From about 300,000 plaques, 17 clones were identified and of these, 7 form overlapping cDNA's that comprise most of the protein-coding sequence (SEQ ID NO:9).

*Developmental and Tissue Distribution of Mouse ptc RNA.* In both the embryonic and adult Northern blots, the *ptc* probe detects a single 8kb message. Further exposure does not reveal any additional minor bands. Developmentally, *ptc* mRNA is present in low levels as early as 7 dpc and becomes quite abundant by 11 and 15 dpc. While the gene is still present at 17 dpc, the Northern blot indicates a clear decrease in the amount of message at this stage. In the adult, *ptc* RNA is present in high amounts in the brain and lung, as well as in moderate amounts in the kidney and liver. Weak signals are detected in heart, spleen, skeletal muscle, and testes.

*In situ Hybridization of Mouse ptc in Whole and Section Embryos.* Northern analysis indicates that *ptc* mRNA is present at 7 dpc, while there is no detectable signal in sections from 7.75 dpc embryos. This discrepancy is explained by the low level of transcription. In contrast, *ptc* is present at high levels along the neural axis of 8.5 dpc embryos. By 11.5 dpc, *ptc* can be detected in the developing lung buds and gut, consistent with its adult Northern profile. In addition, the gene is present at high levels in the ventricular zone of the central nervous system, as well as in the zona limitans of the prosencephalon. *ptc* is also strongly transcribed in the condensing cartilage of 11.5 and 13.5 dpc limb buds, as well as in the ventral portion of the somites, a region which is prospective sclerotome and eventually forms bone in the vertebral column. *ptc* is present in a wide range of tissues from endodermal, mesodermal and ectodermal origin supporting its fundamental role in embryonic development.

*Isolation of the Human ptc Gene.* To isolate human *ptc* (*hptc*), 2 x 10<sup>5</sup> plaques from a human lung cDNA library (HL3022a, Clonotech) were screened with a 1kbp mouse *ptc* fragment, M2-2. Filters were hybridized overnight at reduced stringency (60° C in 5X SSC, 10% dextran sulfate, 5X Denhardt's, 0.2 mg/ml sonicated salmon sperm DNA, and 0.5% SDS). Two positive plaques (H1 and H2) were isolated, the inserts cloned into pBluescript, and upon sequencing, both contained sequence highly similar to the mouse *ptc* homolog. To

isolate the 5' end, an additional 6 x 10<sup>5</sup> plaques were screened in duplicate with M2-3 EcoRI and M2-3 Xho I (containing 5' untranslated sequence of mouse *ptc*) probes. Ten plaques were purified and of these, inserts were subcloned into pBluescript. To obtain the full coding sequence, H2 was fully and H14, H20, and H21 were partially sequenced. The 5.1kbp of human *ptc* sequence (SEQ ID NO:18) contains an open reading frame of 1447 amino acids (SEQ ID NO:19) that is 96% identical and 98% similar to mouse *ptc*. The 5' and 3' untranslated sequences of human *ptc* (SEQ ID NO:18) are also highly similar to mouse *ptc* (SEQ ID NO:19) suggesting conserved regulatory sequence.

*Comparison of Mouse, Human, Fly and Butterfly Sequences.* The deduced mouse *ptc* protein sequence (SEQ ID NO:10) has about 38% identical amino acids to fly *ptc* over about 1,200 amino acids. This amount of conservation is dispersed through much of the protein excepting the C-terminal region. The mouse protein also has a 50 amino acid insert relative to the fly protein. Based on the sequence conservation of *ptc* and the functional conservation of *hedgehog* between fly and mouse, one concludes that *ptc* functions similarly in the two organisms. A comparison of the amino acid sequences of mouse (*mptc*) (SEQ ID NO:10), human (*hptc*) (SEQ ID NO:19), butterfly (*bptc*) (SEQ ID NO:4) and *drosophila* (*ptc*) (SEQ ID NO:6) is shown in the following table.

#### ALIGNMENT OF HUMAN, MOUSE, FLY, AND BUTTERFLY *PTC* HOMOLOGS

20	HPTC	MASAGNAAEPQDR--GGGGSGCIGAPGRPAGGGRRRRTGGLRRAAAPDRDYLHRPSYCD
	MPTC	MASAGNAA-----GALGRQAGGGRRRRTGGPHRA-APDRDYLHRPSYCD
	PTC	M-----DRDSLPRVPDTHGD--VVDE-----KLFSDL-----YI-RTSWVDA
	BPTC	MVAPDSEAPSNPRITAAHESPCATEA-----RHSADL-----YI-RTSWVDA
		* . . . . . * . . . . . *
25	HPTC	AFALEQISKGKATGRKAPLWLRKAFQRLLFKLGCIYQKNCGKFLVVGLLIFGAFVGLKA
	MPTC	AFALEQISKGKATGRKAPLWLRKAFQRLLFKLGCIYQKNCGKFLVVGLLIFGAFVGLKA
	PTC	QVALDQIDKGKARGSRITAIYLSVVFQSHLETGSSVQKHAGKVLFFVAILVLSTFCVGLKS
	BPTC	ALALSELEKGNIEGGRTSLWIRAWLQEQFLILGCFLQGDAGKVLFFVAILVLSTFCVGLKS
30		** . . . . . * . . . . . * . . . . . * . . . . . * . . . . . *
	HPTC	ANLETNVEELWVEVGGRVSRELNYTRQKIGEEAMFNPQLMIQTPKEEGANVLTTEALLQH
	MPTC	ANLETNVEELWVEVGGRVSRELNYTRQKIGEEAMFNPQLMIQTPKEEGANVLTTEALLQH
	PTC	AQIHSKVHQLWIEGGRLEAELAYTQKTIGEDSATHQLLIQTTHDPNASVLHPQALLAH
35	BPTC	AQIHTRVDQLWVQEGGRLEAELKYTAQALGEADSSSTHQLVIQTAKDPDVSLLHPGALLEH
		* . . . . . * . . . . . * . . . . . * . . . . . * . . . . . *
	HPTC	LDSALQASRVHVYMYNRQWKLEHLCYKSGELITET-GYMDQIIIEYLYPCLIIITPLDCFWE
	MPTC	LDSALQASRVHVYMYNRQWKLEHLCYKSGELITET-GYMDQIIIEYLYPCLIIITPLDCFWE
40	PTC	LEVLVKATAVKVHLYDTEWGLRDMCNPSTPSFEGIIYIEQILRHLIPCSIIITPLDCFWE
	BPTC	LKVVAATRVTVHMYDIEWRLKDLCSYSPSIDFEGYHHIESIIDNVIPCAIITPLDCFWE
		* . . . . . * . . . . . * . . . . . * . . . . . *
	HPTC	GAKLQSGTAYLLGKPPLR---WTNFDPLEFLEELK-----KINYQVDSWEEMLNKAEV
45	MPTC	GAKLQSGTAYLLGKPPLR---WTNFDPLEFLEELK-----KINYQVDSWEEMLNKAEV
	PTC	GSQLL-GPESAVVIPGLNQRLRLWTTLNPASVMQYMKQKMSEEKISFDFETVEQYMKRAAI
	BPTC	GSKLL-GPDYPIYVPHLKHKLQWTHLNPLEVVEEVK-KL---KFQFPLSTIEAYMKRAGI
		* . . . . . * . . . . . * . . . . . * . . . . . *

Variable	Mean	SD	Min	Max
Age	34.5	10.2	18	65
Gender	Male	Female		
Marital Status	Married	Single		
Education	High School	College		
Occupation	Manager	Worker		
Income	\$30,000	\$40,000		
Health Status	Good	Fair		
Stress Level	Low	High		
Life Satisfaction	High	Low		
Work-Life Balance	Good	Poor		
Family Support	Strong	Weak		
Community Involvement	Active	Passive		
Religious Beliefs	Religious	Secular		
Political Views	Conservative	Liberal		
Environmental Concerns	High	Low		
Technology Use	Frequent	Infrequent		
Travel Habits	Frequent	Infrequent		
Dietary Preferences	Vegetarian	Non-Vegetarian		
Exercise Routine	Regular	Irregular		
Sleep Patterns	Regular	Irregular		
Substance Use	None	Occasional		
Personal Goals	Clear	Vague		
Resilience	High	Low		
Empathy	High	Low		
Communication Skills	Good	Poor		
Problem Solving	Effective	Ineffective		
Decision Making	Rational	Emotional		
Time Management	Good	Poor		
Organization	High	Low		
Attention Span	Long	Short		
Memory	Good	Poor		
Learning Style	Visual	Auditory		
Creativity	High	Low		
Adaptability	High	Low		
Flexibility	High	Low		
Openness	High	Low		
Conscientiousness	High	Low		
Agreeableness	High	Low		
Neuroticism	Low	High		
Extraversion	High	Low		
Introversion	Low	High		
Social Skills	Good	Poor		
Interpersonal Relationships	Good	Poor		
Conflict Resolution	Effective	Ineffective		
Leadership Skills	Good	Poor		
Teamwork	Good	Poor		
Networking	Active	Passive		
Public Speaking	Good	Poor		
Writing Skills	Good	Poor		
Reading Habits	Frequent	Infrequent		
Learning New Skills	Willing	Unwilling		
Personal Growth	High	Low		
Self-Reflection	Regular	Irregular		
Goal Setting	Clear	Vague		
Time Management	Good	Poor		
Organization	High	Low		
Attention Span	Long	Short		
Memory	Good	Poor		
Learning Style	Visual	Auditory		
Creativity	High	Low		
Adaptability	High	Low		
Flexibility	High	Low		
Openness	High	Low		
Conscientiousness	High	Low		
Agreeableness	High	Low		
Neuroticism	Low	High		
Extraversion	High	Low		
Introversion	Low	High		
Social Skills	Good	Poor		
Interpersonal Relationships	Good	Poor		
Conflict Resolution	Effective	Ineffective		
Leadership Skills	Good	Poor		
Teamwork	Good	Poor		
Networking	Active	Passive		
Public Speaking	Good	Poor		
Writing Skills	Good	Poor		
Reading Habits	Frequent	Infrequent		
Learning New Skills	Willing	Unwilling		
Personal Growth	High	Low		
Self-Reflection	Regular	Irregular		
Goal Setting	Clear	Vague		

	HPTC	VLAYKLLVQTGSRDKPIDISQLTK-QRLVDADGI INPSAFYIYLTAWVSNDPVAYAASQA
	MPTC	VLAYKLLVQTGSRDKPIDISQLTK-QRLVDADGI INPSAFYIYLTAWVSNDPVAYAASQA
	PTC	ILAYKLIVQTHGVNDPNVDKELVLT-NRLVNSDGI INQRAFYNLSAWATNDVFAYGASQG
5	BPTC	ILAYKLMVQTHGVNDNPIDKSLITAGHRLVDKDGII NPKAFYNLSAWATNDALAYGASQG
	HPTC	NIRPHRPEWVHDKADYMPETRLRIPAAEPIEYAQFPFYLNGLRDTSDFVEAIEKVRTICS
	MPTC	NIRPHRPEWVHDKADYMPETRLRIPAAEPIEYAQFPFYLNGLRDTSDFVEAIEKVRVICN
	PTC	KLYPEPRQYFHHQPNY-----DLKIPKSLPLVYAQMPPFYHGLTDTDSQIKTLIGHIRDLSV
10	BPTC	NLKPQPQRWHSPEVD-----HLEIKKSSPLIYTLQPLFYLSGLSDTDSIKTLIRSVRDLCL
	HPTC	NYTSLGLSSYPNGYPFLFWEQYIGLPHWLLLFISVVLACTFLVCVAVLLNPWTAGIIVMV
	MPTC	NYTSLGLSSYPNGYPFLFWEQYISLRHWLLLSISVVLACTFLVCVAVLLNPWTAGIIVMV
15	PTC	KYEGFGLPNYPSGIPFIFWEQYMTLRSSLAMILACVLLAALVLVSLLLSVWAAVLVILS
	BPTC	KYEAKGLPNFPSGIPFLFWEQYLYLRTSLLLALACALGAVFIAVMVLLLNAAVAVLVTLA
	HPTC	LALMTVELFGMMGLIGIKLSAVPVVILIASVGIGVEFTVHVALAFLTAIGDKNRRRAVLAL
20	MPTC	LALMTVELFGMMGLIGIKLSAVPVVILIASVGIGVEFTVHVALAFLTAIGDKNHRAVLAL
	PTC	VLASLAQIFGAMTLLGIKLSAIPAVILILSVGMMLCFNVLISLGFMTSVGNRQRRVQLSM
	BPTC	LATLVLQLLGVMALLGVKLSAMPVLLVLAIGRGVHFTVHLCGLGFVTSIGCKRRRASLAL
25	HPTC	EHMFAPVLDGAVSTLLGVLMLAGSEFDFIVRYFFAVLAILTILGVNLGLVLLPVLLSFFG
	MPTC	EHMFAPVLDGAVSTLLGVLMLAGSEFDFIVRYFFAVLAILTVLGVNLGLVLLPVLLSFFG
	PTC	QMSLGPLVHGMILTSGVAVFMLTSTSPFEFVIPHFCWLLLVLCVAGCNSLLVFPILLSMVG
	BPTC	ESVLAPVVHGAALAAALASMLA.ASEFGFVARLFLRLLLALVFLGLIDGLLFFPIVLSILO
30		
	HPTC	PYPEVSPANGLNRLPTPSPEPPPSVVRFAMPPGHTHSGSDSSDSEYSSQTTVSGLSE-EL
	MPTC	PCPEVSPANGLNRLPTPSPEPPPSVVRFAVPPGHTNNGSDSSDSEYSSQTTVSGISE-EL
	PTC	PEAELVPLEHPDRISTPSPLPVRSSKRSKSYVVGSRSSRGSCQKSHHHHHKDLNDPSL
	BPTC	PAAEVRPIEHPERLSTPSPKCSPIHPRKSSSSSGGDKSSRTS--KSAPRPC-----APSL
35		
	HPTC	RHYEAQQGAGGPAHQVIVEATENPVFAHSTVVHPESRHHPPSNPRQQPHLDSGSLPPGRQ
	MPTC	RQYEAQQGAGGPAHQVIVEATENPVFARSTVVHPDSPHQPLTPRQQPHLDSGSLSPGRQ
	PTC	TTITEEPQSWKSSNSSIQMPNDWTYQPREQ--RPASYAAPPPAYHKAAAQQHHQHQPPT
40	BPTC	TTITEEPSSWHSSAHSVQSSMQSIVVQPEVVVETTTYNNGSDSASGRSTPTKSSHGGAITT
	HPTC	GQQPRRDPPREGLWPPLYRPRRDAFEISTEGHSGPSNRARWGPRGARSHNPPNPASTAMG
	MPTC	GQQPRRDPPREGLWPPPYRPRRDAFEISTEGHSGPSNRDRSGPRGARSHNPRNPTSTAMG
45	PTC	TPPPPFPTA-----YPPQLSIVVQPEVTVETTHS-----DS
	BPTC	TKVTATANIKVEVVTSPDRKSRRSYHYDRRDRDRDRDRDRDRDRDRDRDRDRDRDRDRDR
	HPTC	SSVPGYCQPITTTVTASASVTVAVHPPVPVPGPRNPRGGLCPGY---PETDHGLFEDPHVP
50	MPTC	SSVPSYCQPITTTVTASASVTVAVHPP--PGPRNPRGGPCPGYESYPETDHGVFEDPHVP
	PTC	NT-----TKVTATANIKVELAMP-----GPAVRS---YNFTS-----
	BPTC	DR-----DRERSRERDRP.DRYRD-----EPDHPA---SPRENGRDSGHE-----
55	HPTC	FHVRCERRDSKVEVIELQDVECEERPRGSSSN
	MPTC	FHVRCERRDSKVEVIELQDVECEERPWGSSSN
	PTC	-----
	BPTC	-----SDSSRH

The identity of ten other clones recovered from the mouse library is not determined. These cDNAs cross-hybridize with mouse *ptc* sequence, while differing as to their restriction maps. These genes encode a family of proteins related to the patched protein. Alignment of the human and mouse nucleotide sequences, which includes coding and noncoding sequence, reveals 89% identity.

*Radiation hybrid mapping of the human ptc gene.* Oligonucleotide primers and conditions for specifically amplifying a portion of the human *ptc* gene from genomic DNA by the polymerase chain reaction were developed. This marker was designated STS SHGC-8725. It generates an amplification product of 196 bp, which is observed by agarose gel electrophoresis when human DNA is used as a template, but not when rodent DNA is used. Samples were scored in duplicate for the presence or absence of the 196 bp product in 83 radiation hybrid DNA samples from the Stanford G3 Radiation Hybrid Panel (purchased from Research Genetics, Inc.) By comparison of the pattern of G3 panel scores for those with a series of Genethon meiotic linkage 5 markers, it was determined that the human *ptc* gene had a two point lod score of 1,000 with the meiotic marker D9S287, based on no radiation breaks being observed between the gene and the marker in 83 hybrid cell lines. These results indicate that the *ptc* gene lies within 50-100 kb of the marker. Subsequent physical mapping in YAC and BAC clones confirmed this close linkage estimate. Detailed map information can be obtained from <http://www.shgc.stanford.edu>.

*Analysis of BCNS mutations.* The basal cell nevus syndrome has been mapped to the same region of chromosome 9q as was found for *ptc*. An initial screen of EcoRI digested DNA from probands of 84 BCNS kindreds did not reveal major rearrangements of the *ptc* gene, and so screening was performed for more subtle sequence abnormalities. Using vectorette PCR, by the method according to Riley *et al.* (1990) N.A.R. 18:2887-2890, on a BAC that contains genomic DNA for the entire coding region of *ptc*, the intronic sequence flanking 20 of the 24 exons was determined. Single strand conformational polymorphism analysis of PCR-amplified DNA from normal individuals, BCNS patients and sporadic basal cell carcinomas (BCC) was performed for 20 exons of *ptc* coding sequence. The amplified samples giving abnormal bands on SSCP were then sequenced.

In blood cell DNA from BCNS individuals, four independent sequence changes were found; two in exon 15 and two in exon 10. One 49 year old man was found to have a sequence change in exon 15. His affected sister and daughter have the same alteration, but three unaffected relatives do not. His blood cell DNA has an insertion of 9 base pairs at nucleotide 2445 of the coding sequence, resulting in the insertion of three amino acids (PNI) after amino acid 815. Because the normal sequence preceding the insertion is also PNI, a direct repeat has been formed.

The second case of an exon 15 change is an 18 year old woman who developed jaw cysts at age 9 and BCCs at age 6. The developmental effects together with the BCCs indicate that she has BCNS, although none of her relatives are known to have the syndrome. Her blood cell DNA has a deletion of 11 bp, removing the sequence ATATCCAGCAC at nucleotides 2441 to 2452 of the coding sequence. In addition, nucleotide 2452 is changed from a T to an A. The deletion results in a frameshift that is predicted to truncate the protein after amino acid 813 with the addition of 9 amino acids. The predicted mutant protein is truncated after the seventh transmembrane domain. In *Drosophila*, a *ptc* protein that is truncated after the sixth transmembrane domain is inactive when ectopically expressed, in contrast to the full-length protein, suggesting that the human protein is inactivated by the exon 15 sequence change. The patient with this mutation is the first affected family member, since her parents, age 48 and 50, have neither BCCs nor other signs of the BCNS- DNA from both parents' genes have the normal nucleotide sequence for exon 15, indicating that the alteration in exon 15 arose in the same generation as did the BCNS phenotype. Hence her disease is the result of a new mutation. This sequence change is not detected in 84 control chromosomes.

*Analysis of sporadic basal cell carcinomas.* To determine whether *ptc* is also involved in BCCs that are not associated with the BCNS or germline changes, DNA was examined from 12 sporadic BCCs. Three alterations were found in these tumors. In one tumor, a C to T transition in exon 3 at nucleotide 523 of the coding sequence changes a highly conserved leucine to phenylalanine at residue 175 in the first putative extracellular loop domain. Blood cell DNA from the same individual does not have the alteration, suggesting that it arose somatically in the tumor. SSCP was used to examine exon 3 DNA from 60 individuals who do not have BCNS, and found no changes from the normal sequence. Two other sporadic BCCs have deletions encompassing exon 9 but not extending to exon 8.

The existence of sporadic and hereditary forms of BCCs is reminiscent of the characteristics of the two forms of retinoblastoma. This parallel, and the frequent deletion in tumors of the copy of chromosome 9q predicted by linkage to carry the wild-type allele, demonstrates that the human *ptc* is a tumor suppressor gene. *ptc* represses a variety of genes, including growth factors, during *Drosophila* development and may have the same effect in human skin. The often reported large body size of BCNS patients also could be due to reduced *ptc* function, perhaps due to loss of control of growth factors. The C to T transition identified in *ptc* in the sporadic BCC is also a common genetic change in the *p53* gene in BCC and is consistent with the role of sunlight in causing these tumors. By contrast, the inherited deletion and insertion mutations identified in BCNS patients, as expected, are not those characteristic of ultraviolet mutagenesis.

The identification of the *ptc* mutations as a cause of BCNS links a large body of developmental genetic information to this important human disease. In embryos lacking *ptc* function part of each body segment is transformed into an anterior-posterior mirror-image duplication of another part. The patterning changes in *ptc* mutants are due in part to  
5 derepression of another segment polarity gene, *wingless*, a homolog of the vertebrate Wnt genes that encodes secreted signaling proteins. In normal embryonic development, *ptc* repression of *wg* is relieved by the Hh signaling protein, which emanates from adjacent cells in the posterior part of each segment. The resulting localized *wg* expression in each segment primordium organizes the pattern of bristles on the surface of the animal. The *ptc* gene  
10 inactivates its own transcription, while Hh signaling induces *ptc* transcription.

In flies two other proteins work together with Hh to activate target genes: the ser/thr kinase *fused* and the zinc finger protein encoded by *cubitus interruptus*. Negative regulators working together with *ptc* to repress targets are *protein kinase A* and *costal2*. Thus,  
15 mutations that inactivate human versions of *protein kinase A* or *costal2*, or that cause excessive activity of human *hh*, *gli*, or a *fused* homolog, may modify the BCNS phenotype and be important in tumorigenesis.

In accordance with the subject invention, mammalian patched genes, including the mouse and human genes, are provided, which can serve many purposes. Mutations in the gene are found in patients with basal cell nevus syndrome, and in sporadic basal cell  
20 carcinomas. The autosomal dominant inheritance of BCNS indicates that *patched* is a tumor suppressor gene. The patched protein may be used in a screening for agonists and antagonists, and for assaying for the transcription of *ptc* mRNA. The protein or fragments thereof may be used to produce antibodies specific for the protein or specific epitopes of the protein. In addition, the gene may be employed for investigating embryonic development, by  
25 screening fetal tissue, preparing transgenic animals to serve as models, and the like.

As described above, patients with basal cell nevus syndrome have a high incidence of multiple basal cell carcinomas, medulloblastomas, and meningiomas. Because somatic *ptc* mutations have been found in sporadic basal cell carcinomas, we have screened for *ptc* mutations in several types of sporadic extracutaneous tumors. We found that 2 of 14 sporadic  
30 medulloblastomas bear somatic nonsense mutations in one copy of the gene and also deletion of the other copy. In addition, we identified mis-sense mutations in *ptc* in two of seven breast carcinomas, one of nine meningiomas, and one colon cancer cell line. No *ptc* gene mutations were detected in 10 primary colon carcinomas and eighteen bladder carcinomas.

BCNS<sup>3</sup> (OMIM #109400) is a rare autosomal dominant disease with diverse  
35 phenotypic abnormalities, both tumorous (BCCs, medulloblastomas, and meningiomas) and developmental (misshapen ribs, spina bifida occulta, and skull abnormalities; Gorlin, R.J. (1987) Medicine 66:98-113). The BCNS gene was mapped to chromosome 9q22.3 by

linkage analysis of BCNS families and by LOH analysis in sporadic BCCs (Gallani, M.R. *et al.* (1992) Cell 69:111-117). LOH in sporadic medulloblastomas has been reported in the same chromosome region (Schofield, D. *et al.* (1995) Am J Pathol 146:472-480). Recently, the human homologue of the *Drosophila patched* (PTCII) gene has been mapped to the BCNS region (Hahn, H. *et al.* (1996) Cell 85:841-851; Johnson, R.L. *et al.* (1996) Science 272:1668-1671; Gallani, M.R. *et al.* (1996) Nat Genet 14:78-81; Xie, J. *et al.* (1997) Genes Chromosomes Cancer 18:305-309), and mutations in this gene have been found in the blood DNA of BCNS patients and in the DNA of sporadic BCCs (Hahn, H. *et al.*, *supra*; Johnson, R.L. *et al.*, *supra*; Gallani, M.R. *et al.*, *supra*; and Chidambaram, A. *et al.* (1996) Cancer Res 36:4599-4601). *ptc* appears to function as a tumor suppressor gene; inactivation abrogates its normal inhibition of the hedgehog signaling pathway. Because of the wide variety of tumors in patients with the BCNS and wide tissue distribution of *ptc* gene expression, we have begun screening for *ptc* gene mutations in several types of human cancers, especially those present in increased numbers in BCNS patients (medulloblastomas), those in tissues derived embryologically from epidermis (breast carcinomas) and those with chromosome 9q LOG (bladder carcinomas; see Cairns, P. *et al.* (1993) Cancer Res 53:1230-1232; and Sidransky, D. *et al.* (1997) NEJM 326:737-740).

To further study the roles of *ptc* in development and in tumorigenesis, we have constructed mice lacking *ptc* function. By homologous recombination, part of *ptc* exon 1 (including the putative start codon) and all of exon 2 were replaced with *lacZ* and a neomycin resistance gene (Fig. 3) (DNA from the *ptc* genomic locus was isolated from a 129SV genomic phage library [Stratagene]. Exons 1-15 of human *PTC (1)* were mapped by PCR and sequencing. The 3' arm of homology was a 3.5 kb EcoRI-BamHI fragment from the second intron that gained a BamHI site from pBSII [Stratagene] and was cloned into the BamHI site of pPNT [Tybulewicz, *et al.* (1991) Cell 65:1153]. A cassette containing the gene for nuclear localized b-galactosidase, followed by the mP1 intron and polyA tail was excised from pNLacF [Mercer, *et al.* (1991) Neuron 7:703] and cloned into the Xho I site of pPNT using Xho I and Sal I linkers. The 5' arm of homology was a 6.5 kb Xho I to Nru I fragment that was cloned into the Xho I site upstream of *lacZ* via a Sal I linker. The Nru I site is in the first *ptc* exon. The resulting plasmid, KO1, was linearized with Xho I and electroporated into RI ES cells that were subjected to double selection and analyzed by Southern blot [Joyner, A.L. Gene Targeting: A Practical Approach. Oxford University Press, New York, 1993, pp.33-61]. Targeted clones were expanded and used for injection into C57Bl/6 blastocysts [Hogan, B. *et al.* Manipulating the Mouse Embryo: A Laboratory Manual Cold Spring Harbor Laboratory Press, Cold Spring Harbor, 1994, pp.196-204]. Protein made from any alternative ATG would lack the first proposed transmembrane domain, flipping the orientation of the protein in the membrane. Three independent ES clones were used to make chimeras that were bred to B6D2F1 animals to generate

heterozygous mice on a mixed background. Interbreeding of heterozygotes produced no homozygous animals among 202 offspring examined. Analysis of embryos from timed matings suggested that *ptc*<sup>-/-</sup> embryos die between embryonic day (E) 9.0 and E10.5, with the first gross phenotypes appearing by E8. In *ptc*<sup>-/-</sup> embryos, the neural tube failed to close completely and was overgrown in the head folds, hindbrain and spinal cord (Fig. 4, A to C). Embryonic lethality may have been due to abnormal development of the heart (Fig. 4B), which never beats.

In flies Ptc protein inhibits *ptc* transcription. By inhibiting Ptc function, Hh increases production of Ptc which may then bind available Hh and limit the range or duration of effective Hh signal (Y. Chen and G. Struhl, (1996) Cell 87:553). Hh signaling also post-transcriptionally regulates the zinc finger protein cubitus interruptus (*ci*) (C. K. Motzny and R. Holmgren, (1996) Mech Dev 52:137; Domínguez, *et al.* (1996) Science 272:1621; Hepker, *et al.* (1997) Development 124:549; Aza-Blanc, *et al.*, (1997) Cell 89:1043). In vertebrates, Sonic hedgehog (Shh) signaling induces transcription of both *ptc* and a *ci* homolog, *Gli* (Goodrich, *et al.* (1996) Genes Devel. 10:301; Marigo, *et al.* (1996) Development 122:1225; Concordet, *et al.*, (1996) Development 122:2835; Marigo, *et al.* (1996) Dev. Biol. 180:273). Derepression of *ptc* and *Gli* in *ptc*<sup>-/-</sup> mice should therefore reveal where Ptc is normally active.

*ptc* and *Gli* expression was greatly increased in *ptc*<sup>-/-</sup> embryos. In *ptc*<sup>+/-</sup> mice expression of the *lacZ* gene fused to the first *ptc* exon during targeting accurately reported the pattern of *ptc* transcription (Fig. 4, C and D). In *ptc*<sup>-/-</sup> embryos expression of *ptc-lacZ* was extensively derepressed starting at about E8.0 in the anterior neural tube and spreading posteriorly by E8.75 (Fig. 4, C and E). Derepression was germ layer-specific: both *ptc-lacZ* and *Gli* were expressed throughout the ectoderm and mesoderm, but not in the endoderm (Fig. 4, D to G). *ptc* expression may be excluded from the endoderm in order to avoid interfering with Shh signaling from the endoderm to the mesoderm (Roberts *et al.*, (1995) Development 121:3163). A differential requirement for Ptc may distinguish the germ layers.

As revealed by *ptc* mutants, an early site of Ptc activity is the neural tube, where Shh and Ptc act antagonistically to determine cell fates. Shh induces the floor plate and motor neurons in the ventral neural tube (Echelard *et al.*, (1993) Cell 75:1417; Roelink *et al.*, (1994) Cell 76:761; Roelink *et al.*, (1995) Cell 81:445-455). These cell types fail to form in *Shh* mutants (Chiang *et al.*, (1996) Nature 383:407). High levels of Shh produced by the notochord may induce floor plate by completely inactivating Ptc (Echelard *et al.*, (1993) *supra*; Roelink *et al.*, (1994) *supra*; Roelink *et al.*, (1995) *supra*). If so, elimination of *ptc* function might cause floor plate differentiation throughout the neural tube. Prospective floor plate cells transcribe the forkhead transcription factor *HNF3b* first and then *Shh* itself (Echelard *et al.*, (1993) *supra*; Roelink *et al.*, (1994) *supra*; Roelink *et al.*, (1995) *supra*). In

E8.5 *ptc* mutants, transcription of *HNF3b* and *Shh* was expanded dorsally (Fig. 5, A to C). Ectopic *Shh* expression was most extensive in the anterior, where transcripts could be detected throughout the neurepithelium (Fig. 5, B and C). Cells in this region were in a single layer with basal nuclei, like floor plate cells that are normally restricted to the ventral midline (Fig. 5, D and E). Expression of the intermediate neural tube marker *Pax6* (C. Walther and P. Gruss, (1991) Development 113:1435) was completely absent from *ptc* mutant embryos, suggesting that only ventral, and not ventrolateral, cell fates are specified (Fig. 5, F and G).

Dorsalizing signals from the surface ectoderm (Dickinson, *et al.* (1995) Development 121:2099; Liem, *et al.* (1995) Cell 82:969) could confer dorsal cell fates even in the absence of *ptc* function. In E8-E9 *ptc* homozygotes the dorsal neural tube marker *Pax3* was not expressed in the anterior neural tube, but was transcribed in a very small region at the dorsal-most edge of the posterior neural tube (Fig. 5, H to J). In addition *erb-b3* transcription, which marks migratory neural crest cells (Fig. 5K) (H. U. Wang and D. J. Anderson, (1997) Neuron 18:383), was not detected in the somites of *ptc* mutants (Fig. 5L). We conclude that only limited dorsal fate determination occurs in the absence of *ptc*. BMP signals maintain dorsal gene expression (Dickinson, *et al.* (1995) *supra*; Liem, *et al.* (1995) *supra*), so either *ptc* is required for BMPs to work or BMP signaling is ineffective in most cells expressing *Shh* targets.

Ventralization of the neural tube in *ptc* mutants occurred without affecting cell identity along the rostrocaudal axis. In *ptc*<sup>-/-</sup> embryos, cells in the anterior neural tube expressed the forebrain marker *Nkx2.1* (Shimamura, *et al.* (1995) Development 121:3923) and cells in the spinal cord transcribed low levels of *hoxb1* (Wilkinson, *et al.* (1989) Nature 341:405) (Fig. 5, M and N). *hoxb1* was not transcribed in the fourth rhombomere of *ptc* mutants (Fig. 5, N). This may reflect a transformation of hindbrain cells to floor plate, since *hoxb1* is excluded from the midline of wild-type embryos. Conversely, in the anterior, *Nkx2.1* expression was expanded dorsally in mutants compared to wild-type embryos (Fig. 5, M).

*ptc*<sup>+/-</sup> mice had phenotypes similar to those of BCNS patients: they were larger than their wild-type littermates [ $30.72 \pm 3.83$  (average  $\pm$  SD; n=29) vs.  $26.54 \pm 2.51$  (n=39) at 2-3 months;  $P=0.000001$ ], a small fraction (3 of 389 mice examined) had hindlimb defects such as extra digits or syndactyly (Fig. 6A) or obvious soft tissue tumors (1 of 243) and many developed brain tumors (see below).

Of 243 *ptc*<sup>+/-</sup> mice which were between the ages of 2 and 9 months and were not sacrificed for other studies, 18 died or were euthanized because of sickness. No wild-type littermates died. Ten of the affected heterozygotes were autopsied and eight were found to have large growths in the cerebellum that resembled medulloblastomas (Fig. 6, B and C).

Human medulloblastomas are believed to arise from a "primitive neuroectodermal" cell type (J.P. Provias and L. E. Becker, (1996) J Neurooncol 29:35). They are most common in children, can be metastatic or non-metastatic, and can have glial and neuronal properties. The histology of tumors from *ptc*<sup>+/-</sup> mice was similar to that of human medulloblastoma: tumor cells were small, with dark carrot-shaped nuclei and little cytoplasm (Fig. 6, D and E), and although a subset expressed neurofilament protein and synaptophysin (Fig. 6F) (For immunostaining, two tumors were fixed and embedded in paraffin. Tissue sections (4 mm) were cleared and dehydrated, treated with 3% hydrogen peroxide and then with a dilution of 1:10 normal rabbit serum (Vector Laboratories). Anti-synaptophysin (Boehringer-Mannheim) was used at a dilution of 1:5 and anti-neurofilament protein (Dako) at 1:50. Antibody binding was visualized with a peroxidase Vectastain Elite ABC kit (Vector Laboratories). Nuclei were counterstained with hematoxylin. Like anti-synaptophysin, anti-neurofilament staining appeared in processes of the tumor cells.), the majority of cells appeared undifferentiated. Of the two autopsied animals without apparent medulloblastomas, one had a large tumor growing out of its rib muscle and the other died for unknown reasons. Medulloblastomas and soft tissue tumors were also observed in *ptc*<sup>+/-</sup> mice maintained on an inbred 129SV background: 6 of 27 had obvious medulloblastomas; 2 of 27 had tumors in the muscle of their leg; and 3 of 27 died but were not examined.

The *ptc* and *Gli* genes were strongly transcribed in the brain tumors but not in surrounding tissue (Fig. 7, A and B; n = 3 of 3 tumors examined). There was no detectable increase in *Shh* expression (Fig. 7C). To assess the incidence of medulloblastomas, brains from 47 asymptomatic *ptc*<sup>+/-</sup> mice were randomly collected and stained with X-gal. Nine brains contained medulloblastomas that were easily recognized by their disorganized morphology and intense *ptc-lacZ* expression (Fig. 7D). Medulloblastomas were observed in 7 of 23 (30.4%) *ptc*<sup>+/-</sup> mice at 12 to 25 weeks of age, 1 of 12 (8.3%) mice at 9 to 10 weeks and 1 of 12 (8.3%) mice at 5 weeks. Tumors can therefore arise as early as 5 weeks postnatally, but they increase in severity and frequency as the animal ages.

We looked for changes in *ptc-lacZ* expression that might reflect early stages of tumorigenesis. At all stages examined, about half of the animals [50% at 5 to 10 weeks (n=24), 56.5% at 12 to 25 weeks (n=23)] exhibited regions of increased X-gal staining on the surface of the cerebellum (Fig. 7E). These regions were usually lateral and often extended down into the fissures separating the folia (Fig. 7, E and F). The mouse medulloblastomas may arise from these cells, which are superficial to the molecular layer of the cerebellum (Fig. 7F). During fetal development, prospective cerebellar granule cells proliferate in the external granule layer (EGL), the outermost layer of the cerebellum. Granule cells then leave and migrate past the Purkinje cells to form the internal granule cell layer of the adult animal, gradually depleting the EGL. The remnants of the fetal EGL have been proposed to be a source of human medulloblastoma progenitors, a hypothesis consistent with the higher

frequency of these tumors in children (L. Stevenson and F. Echlin, (1934) Arch. Neurol. Psychiat. 31:93; Kadin, *et al.* (1970) J Neuropathol Exp Neurol 29:583).

The abundance of cerebellar *ptc* transcripts was reduced by about 50% in the *ptc*<sup>+/-</sup> mice compared to wild-type littermates (Fig. 7G). This reduction could lead to ectopic expression of Shh target genes and to uncontrolled cell proliferation. Brain tumors might arise from *Ptc* haploinsufficiency alone, from additional mutations in the second *ptc* allele, or from a combination of *ptc* mutations with mutations in other tumor suppressor loci. We have not observed BCCs in *ptc*<sup>+/-</sup> mice, perhaps because somatic inactivation of the second *ptc* gene is required as it is in human BCCs.

Our analysis has revealed that *Ptc* controls growth and pattern formation in early neural development and in the adult cerebellum. Autoregulation of *ptc* occurs in vertebrates as it does in flies, and the balance between Hh and *Ptc* activities appears critical for normal development. The importance of *Ptc* dosage is emphasized by the phenotype of the *ptc*<sup>+/-</sup> mice, which develop a tumor type observed in the corresponding human cancer predisposition syndrome. Medulloblastoma is a common childhood brain tumor and the prognosis remains grim. The Hh/*Ptc* pathway may provide new diagnostic tools and new insights into tumorigenesis that may be directed toward potential therapies.

## Materials and methods

Clinical Materials. Diagnoses of all tumors were confirmed histologically. Cell lines were obtained from the America Type Culture Collection. DNA was extracted from tumors or matched normal tissue (peripheral blood leukocytes or skin) as described (Cogen, P.H. *et al.* (1990) Genomics 8:279-285; and Sambrook, J. *et al.* Molecular Cloning: A Laboratory Manual, Ed. 2, Vol. 2, pp. 9.17 - 9.19, Cold Spring Harbor, NY (1989)).

PCR and Heteroduplex Analysis. PCR amplification and heteroduplex/SSCP analysis were performed as described (Johnson, R.L. *et al.*, *supra*; Spritz, R.A. *et al.* (1992) Am J Hum Genet 51:1058-1065). Primers used and intron/exon boundary sequences of the *ptc* gene were derived as reported previously (Johnson, R.L. *et al.*, *supra*) and are shown in Table 1. Primers for exon 1 and 2 were from Hahn *et al.* (*supra*).

Sequence Analysis. Exon segments exhibiting bands were reamplified and were sequenced directly using the Sequenase sequencing kit according to the protocol recommended by the manufacturer (United States Biochemical Corp.). A second sequencing was performed using independently amplified PCR products to confirm the sequence change. The amplified PCR products from each tumor were also cloned into the plasmid vector pCR 2.1 (InVitrogen), followed by sequence analysis of at least four independent clones. The sequence alteration was confirmed from at least two independent clones. Simplified

amplification of specific allele analysis was performed according to Lei and Hall (Lei, X. and Hall, B.G. (1994) Biotechniques 16:44-45).

Allele Loss Analysis. Microsatellites used for allelic loss analysis were D9S109, DpS119, D9S127, D9S196, and D9S287 described in the CHLC human screening set  
 5 (Research Genetics). A part of the *ptc* intron 1 sequence was tested for polymorphism in a control population and found to be polymorphic in 80% of the samples tested. This microsatellite was used for analysis of *ptc* gene allelic loss in bladder carcinomas. The primer sequences are as follows: forward primer, 5'-CTGAGCAGATTTCCCAGGTC-3'; and reverse primer, 5'-CCTCAGACAGACCTTTCCTC-3'. The PCR cycling for this newly  
 10 isolated marker was 4 min. at 95 C, followed by 30 cycles of 40 s at 95 C, 2 min. at 60 C, and 1 min. at 72 C. PCR products were separated on 6% polyacrylamide gels and exposed to film.

## Results and Discussion

Intronic boundaries were determined for 22 exons of *ptc* by sequencing vectorette  
 15 PCR products derived from BAC 192J22 (Johnson R.L., *supra*; Table 1). Our findings are in agreement with those of Hahn *et al.* (*supra*), except that we find exon 12 is composed of 2 separate exons of 126 and 119 nucleotides. This indicates that *ptc* is composed of 23 coding exons instead of 22. In addition, we find that exons 3, 4, 10, 11, 17, 21, and 23 differ slightly  
 20 in size than reported previously (Hahn *et al.*, *supra*). Of 63 tumors studied, 14 were sporadic medulloblastomas, and 9 were sporadic meningiomas. These 23 tumors were examined for allelic deletions by genotyping of tumor and blood DNA with microsatellite markers that flank the *ptc* gene: D9S119, D9S196, D9S287, D9S127, and D9S109. Four of 14 medulloblastomas had LOH. Two of the medulloblastomas, both of which had LOH, had  
 25 mutations (med34 and med36; see Cogen, P.H. *et al.*, *supra*), which are predicted to result in truncated proteins (Table 2). DNA samples from the blood of these patients lack these mutations, indicating that they both are somatic mutations. med34 also has allelic loss on 17p (Cogen, P.H. *et al.*, *supra*). We were unable to detect *ptc* gene mutations by heteroduplex analysis in the other two medulloblastomas bearing LOH on 9q. The  
 30 pathological features of these two tumors differed in that med34 belongs to the desmoplastic subtype, whereas med36 is of the classic type, indicating that *ptc* mutations in medulloblastomas are not restricted to a specific subtype.

"041915" 457230

TABLE 1 Primers and boundary sequences of *PTCH*

Exon	5' Boundary <sup>a</sup>	Nucleotide position <sup>b</sup>	Exon size	3' boundary <sup>c</sup>	Reading frame <sup>d</sup>	Primers
1	ND <sup>e</sup>	ND	ND	ag <sup>g</sup> GTGAT	ND	
2	ND	202	193	ag <sup>g</sup> GTAAAG	3	
3	TGTCAG <sub>12</sub>	323	190	ag <sup>g</sup> GTAAAG	1	3F GAATTTCGAGTATGTTGCTNITC 3R ACCGCTTACCTGCTGCTC 4F TGCACTAATTTCTTATACACTC 4R TAAAGCACACTACTGGGCTC 5F GAACACCCCACTAGTGTGCTC 5R TGAGTCCCTAGAGAAAGTCACAGACA 6F GGTCTCTTTGATGCTCTGCTC 6R TTTTTCCTCTCCACCTTC 7F GCACGTGATTTTAAACAGGCTC 7R AGGOCATAGATTTCTCTGGG 8F TGGGAATACGTGATGATGCTGCTC 8R CATAACCCAGGCTCTGCTC 9F CATTTGGGCTTTGGGCTC 9R ACCAAACCAACTCCAGCTC 10F TGCCCCCATTTGTTCTGCTC 10R GCACAGCAGATAAATGGCTC 11F GCATCTGGCATGCTTAAAGCTC 11R AAGCTGTGATGCTCCCAAG 12F GACCATGCTCCAGTCCAGCTC 12R COTTCAGATCAGCAAGCTC 13F AGTCTCTGATTTGGGCTC 13R CCATTCTGACCCCAATCAAAAC 14F AAATGGCAGAAATGAAGGCTC 14R CTGATGAACCTCAAGCTCTC 15F GGAAGAGTCACTGCTGCTC 15R CGCCAAAGCCGAAAGCTC 16F AGGCTCTCTGCTGCTGCTC 16R GCTGCTCAAGGCTGCTC 17F GCTCTCAAGGCTGCTGCTC 17R GGAAGGCTCTGCTGCTGCTC 18F GCTCTCAAGCTGCTGCTGCTC 18R GAATTTGACTTCCAGCAAGCTC 19F GCGCCAGTGAACCTGCTGCTC 19R GAGCCAGAGGATGCTGCTC 20F AGCATTACCGGCTGCTGCTC 20R TTGCACACGCTGCTGCTC 21F TTTTCCGCTTCTGCTGCTC 21R GCACAGGAAACGCTGCTC 22F GCAAGTAAATGCAAGGCTC 22R ACTACACGCTGCTGCTC 23F CCTTCTAAGCCGCTGCTC 23R GACACATCAGCTGCTC
4	TATTAG <sub>12</sub>	555	70	ag <sup>g</sup> GTATAT	2	
5	TGACAG <sub>12</sub>	655	92	cc <sup>g</sup> GTAAAT	3	
6	TGACAG <sub>12</sub>	747	199	aa <sup>g</sup> GTGAGT	2	
7	TTT <sub>12</sub> AG <sub>12</sub>	945	122	ag <sup>g</sup> GTAAAG	3	
8	CTGCAG <sub>12</sub>	1058	148	ag <sup>g</sup> GTAAAC	2	
9	CCACAG <sub>12</sub>	1216	132	ag <sup>g</sup> GTAAAG	3	
10	TTCCAG <sub>12</sub>	1348	156	ag <sup>g</sup> GTACTA	3	
11	CTGTAG <sub>12</sub>	1504	99	ag <sup>g</sup> GTAAAT	3	
12	TCCAG <sub>12</sub>	1603	126	ag <sup>g</sup> GTGAGC	3	
13	TCCAG <sub>12</sub>	1729	119	ag <sup>g</sup> GTACAT	3	
14	TTT <sub>12</sub> AG <sub>12</sub>	1848	403	ag <sup>g</sup> GTAAAT	2	
15	TTCCAG <sub>12</sub>	2251	310	ag <sup>g</sup> GTAAAG	3	
16	TCTAG <sub>12</sub>	2561	143	ag <sup>g</sup> GTACTC	1	
17	TTGTAG <sub>12</sub>	2704	184	aa <sup>g</sup> GTAAAT	3	
18	GTCCAG <sub>12</sub>	2888	281	aa <sup>g</sup> GTGAGT	1	
19	CTCCAG <sub>12</sub>	3159	138	ag <sup>g</sup> GTATGG	3	
20	GCACAG <sub>12</sub>	3307	143	ag <sup>g</sup> GTAAAG	3	
21	TCCAG <sub>12</sub>	3450	100	ag <sup>g</sup> GTCAAGT	2	
22	AAATAG <sub>12</sub>	3550	255	aa <sup>g</sup> GTAAAT	3	
23	CTCCAG <sub>12</sub>	3805	341	ag <sup>g</sup> GTGAGT	3	
24	ND	4346	ND	ND		

Consensus sequences for the 5' and 3' exonic boundaries are (T)<sub>12</sub>NCAAG and ag<sup>g</sup>GTAAAGT, respectively (20) (lower case denotes exonic sequence).  
<sup>a</sup>From positions are in reference to the coding sequence of *PTCH* (3) with the beginning ATG as nucleotide 1.  
<sup>b</sup>5' exon boundary begins after the first, second, or third base of the codon of the translational reading frame.  
<sup>c</sup>ND, not determined.

One report (Schofield, D. *et al.*, *supra*) has shown that five medulloblastomas (two BCNS-associated cases and three sporadic cases) bearing LOH on chromosome 9q22.3-q31 are all of the desmoplastic subtype, suggesting LOH on 9q22.3 is histological subtype specific. We feel that the conclusion derived from only five positive tumors is a not strong one because we and others (Raffel, C. *et al.* (1997) *Cancer Res* 57:842-845) have found nondesmoplastic subtypes of medulloblastomas bearing LOH on chromosome 9q22.3. Independently, another group has reported their finding of *ptc* mutations in sporadic medulloblastomas (Raffel, C. *et al.*, *supra*).

A change of T to C at nucleotide 2990 (in exon 18) was identified in DNA from one of nine sporadic meningiomas, causing a predicted change of codon 997 from Ile to Thr (Table 2). The meningioma bearing this mutation also has allelic loss on 9q22.3. Blood cell DNA is heterozygous for this mutation, but DNA from the tumor contains only the mutant sequence. Of 100 normal chromosomes examined, none has this sequence change, suggesting that this mutation is not likely a common polymorphism. This patient is 84 years



and Hall, B.G., *supra*) to examine 50 normal blood DNA samples for the presence of the sequence alteration and found none but the DNA from this cell line to have the mutant allele, suggesting that this mutation also is unlikely to be a common sequence polymorphism. For bladder carcinomas, a newly isolated microsatellite that was derived from intron 1 of the *ptc* gene was used to examine LOH in the tumor. Three primary bladder carcinomas showed LOH at this intragenic locus. With no *ptc* mutations detected in these tumors, we suspect that the LOH in these three bladder carcinomas may reflect the high incidence of while chromosome 9 loss in bladder cancers (Sidransky, D. *et al.*, *supra*). A similar observation has been reported previously (Simoneau, A. R. *et al.* (1996) Cancer Res 56:5039-5043).

We also detected a sequence change in intron 10 in two colon carcinomas, 15-1 and 8-1, an alteration that was reported previously as a splicing mutation (Unden, A.B. *et al.* (1996) Cancer Res 56:4562-4565). Because we found the same sequence change in about 20% of normal control samples, we suggest that this more likely is a nonpathogenic polymorphism. The *ptc* protein is predicted to contain 12 transmembrane domains, two large extracellular loops, and one intracellular loop (Goodrich, L.V. *et al.*, *supra*). Of the six mutations we identified, four are missense mutations. Three mutations lead to amino acid substitutions in the second extracellular loop, and one mutation results in an amino acid change in the intracellular domain.

Our data indicate that somatic inactivation of the *ptc* gene does occur in some sporadic medulloblastomas. In addition, because missense mutations of the *ptc* gene were detected in breast carcinomas, we suspect that defects of the *ptc* function also may be involved in some breast carcinomas, although biochemical evidence is necessary to show how these missense mutations might impair *ptc* function. Of 11 colon cancers and 18 bladder carcinomas examined, we found only one mutation in 1 colon cell line, suggesting that *ptc* gene mutations are relatively uncommon in colon and bladder cancers, although the incidence of chromosome 9 loss in bladder cancers is high (Cairns, P. *et al.*, *supra*).

Published reports of SSCP analysis of tumor DNA identified mutations in the *ptc* gene in only 30% of sporadic BCCs, although chromosome 9q22.3 LOH was reported in more than 50% of these tumors (Gallani, M.R. *et al.*, *supra*). It has been reported that heteroduplex/SSCP analysis of gene mutations is more sensitive than SSCP analysis (Spritz, R.A. *et al.*, *supra*). In our studies, we were able to identify a point mutation in the 310-bp - PCR product from exon 15 using heteroduplex analysis, whereas SSCP analysis failed to reveal this sequence change (Table 2). Therefore, we suspect that there may be more mutations in BCCs than we have found thus far. Analysis of the *ptc* gene in BCNS patients and in sporadic BCCs has identified mutations scattered widely across the gene, and the majority of mutations were predicted to result in truncated proteins (Hahn, H. *et al.*, *supra*; Johnson, R.L. *et al.*, *supra*; Gallani, M.R. *et al.*, *supra*; Chidambaram, A. *et al.*, *supra*;

Unden, A.B. *et al.*, *supra*; Wicking, C. *et al.* (1997) Am J Hum Genet 60:21-26). In our screening, we found two breast carcinomas bearing missense mutations of the *ptc* gene. In one of these two tumors, B349, direct sequencing indicated a deletion of the other copy of the *ptc* gene. Any comparison of mutations in skin cancers versus extracutaneous tumors must  
5 consider the wholly different causes of these mutations; UV light is unique to the skin.

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent o application were specifically and individually indicated to be incorporated by reference.

10 Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

DEPOSITED

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

## (ii) TITLE OF INVENTION:

## (iii) NUMBER OF SEQUENCES: 19

## (iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Foley, Hoag & Eliot LLP  
(B) STREET: One Post Office Square  
(C) CITY: Boston  
(D) STATE: MA  
(E) COUNTRY: US  
(F) ZIP: 02109

## (v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

## (vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:  
(B) FILING DATE:  
(C) CLASSIFICATION:

## (viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Vincent, Matthew P.  
(B) REGISTRATION NUMBER: 36,709  
(C) REFERENCE/DOCKET NUMBER: SUV003.04

## (ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 617-832-1000  
(B) TELEFAX: 617-832-7000

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 736 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

AACNNC NNTN NATGGCACCC CCNCCCAACC TTTNNNCCNN NTAANCAAAA NNCCCCNTTT 60  
NATACCCCT NTAANANTTT TCCACCNNNC NNAAANNCCN CTGNANACNA NGNAAANCCN 120  
TTTTTNAACC CCCCCCACCC GGAATTCCNA NTNNCCNCCC CCAAATTACA ACTCCAGNCC 180

AAAATTNANA NAATTGGTCC TAACCTAACC NATNGTTGTT ACGGTTTCCC CCCCCAAATA 240  
 CATGCACTGG CCCGAACACT TGATCGTTGC CGTTCCAATA AGAATAAATC TGGTCATATT 300  
 AAACAAGCCN AAAGCTTTAC AAAGTGTGT ACAATTAATG GGCGAACACG AACTGTTTGA 360  
 ATTCTGGTCT GGACATTACA AAGTGCACCA CATCGGATGG AACCAGGAGA AGGCCACAAC 420  
 CGTACTGAAC GCCTGGCAGA AGAAGTTCGC ACAGGTTGGT GGTGGCGCA AGGAGTAGAG 480  
 TGAATGGTGG TAATTTTTTG TTGTTCCAGG AGGTGGATCG TCTGACGAAG AGCAAGAAGT 540  
 CGTCGAATTA CATCTTCGTG ACGTTCTCCA CCGCCAATTT GAACAAGATG TTGAAGGAGG 600  
 CGTCGAANAC GGACGTGGTG AAGCTGGGGG TGGTGCTGGG GGTGGCGGCG GTGTACGGGT 660  
 GGGTGGCCCA GTCGGGGCTG GCTGCCTTGG GAGTGCTGGT CTTNGCGNGC TNCNATTTCG 720  
 CCTATAGTNA GNCGTA 736

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 107 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Xaa	Pro	Pro	Pro	Asn	Tyr	Asn	Ser	Xaa	Pro	Lys	Xaa	Xaa	Xaa	Leu	Val	1	5	10	15
Leu	Thr	Pro	Xaa	Val	Val	Thr	Val	Ser	Pro	Pro	Lys	Tyr	Met	His	Trp	20	25	30	
Pro	Glu	His	Leu	Ile	Val	Ala	Val	Pro	Ile	Arg	Ile	Asn	Leu	Val	Ile	35	40	45	
Leu	Asn	Lys	Pro	Lys	Ala	Leu	Gln	Thr	Val	Val	Gln	Leu	Met	Gly	Glu	50	55	60	
His	Glu	Leu	Phe	Glu	Phe	Trp	Ser	Gly	His	Tyr	Lys	Val	His	His	Ile	65	70	75	80
Gly	Trp	Asn	Gln	Glu	Lys	Ala	Thr	Thr	Val	Leu	Asn	Ala	Trp	Gln	Lys	85	90	95	
Lys	Phe	Ala	Gln	Val	Gly	Gly	Trp	Arg	Lys	Glu	100	105							

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 5187 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

J57330" 04T9T99

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GGGTCTGTCA CCCGGAGCCG GAGTCCCCGG CGGCCAGCAG CGTCCTCGCG AGCCGAGCGC	60
CCAGGCGCGC CCGGAGCCCG CGGCGGCGGC GGCAACATGG CCTCGGCTGG TAACGCCGCC	120
GGGGCCCTGG GCAGGCAGGC CGGCGGCGGG AGGCGCAGAC GGACCGGGGG ACCGCACCGC	180
GCCGCGCCGG ACCGGGACTA TCTGCACCGG CCCAGCTACT GCGACGCCGC CTTCGCTCTG	240
GAGCAGATTT CCAAGGGGAA GGCTACTGGC CGGAAAGCGC CGCTGTGGCT GAGAGCGAAG	300
TTTCAGAGAC TCTTATTTAA ACTGGGTTGT TACATTCAA AGAACTGCGG CAAGTTTTTG	360
GTTGTGGGTC TCCTCATATT TGGGGCCTTC GCTGTGGGAT TAAAGGCAGC TAATCTCGAG	420
ACCAACGTGG AGGAGCTGTG GGTGGAAGTT GGTGGACGAG TGAGTCGAGA ATTAAATTAT	480
ACCCGTCAGA AGATAGGAGA AGAGGCTATG TTTAATCCTC AACTCATGAT ACAGACTCCA	540
AAAGAAGAAG GCGCTAATGT TCTGACCACA GAGGCTCTCC TGCAACACCT GGACTCAGCA	600
CTCCAGGCCA GTCGTGTGCA CGTCTACATG TATAACAGGC AATGGAAGTT GGAACATTG	660
TGCTACAAAT CAGGGGAACT TATCACGGAG ACAGGTTACA TGGATCAGAT AATAGAATAC	720
CTTTACCCTT GCTTAATCAT TACACCTTTG GACTGCTTCT GGAAGGGGC AAAGCTACAG	780
TCCGGGACAG CATACTCCT AGGTAAGCCT CCTTTACGGT GGACAACTT TGACCCCTTG	840
GAATTCCTAG AAGAGTTAAA GAAAATAAAC TACCAAGTGG ACAGCTGGGA GGAAATGCTG	900
AATAAAGCCG AAGTTGGCCA TGGGTACATG GACCGGCCTT GCCTCAACCC AGCCGACCCA	960
GATTGCCCTG CCACAGCCCC TAACAAAAAT TCAACCAAAC CTCTTGATGT GGCCCTTGTT	1020
TTGAATGGTG GATGTCAAGG TTTATCCAGG AAGTATATGC ATTGGCAGGA GGAGTTGATT	1080
GTGGGTGGTA CCGTCAAGAA TGCCACTGGA AACTTGTCA GCGCTCACGC CCTGCAAACC	1140
ATGTTCCAGT TAATGACTCC CAAGCAAATG TATGAACACT TCAGGGGCTA CGACTATGTC	1200
TCTCACATCA ACTGGAATGA AGACAGGGCA GCCGCCATCC TGGAGGCCTG GCAGAGGACT	1260
TACGTGGAGG TGGTTCATCA AAGTGTGCGC CCAAACCTCA CTCAAAGGT GCTTCCCTTC	1320
ACAACCACGA CCCTGGACGA CATCCTAAAA TCCTTCTCTG ATGTCAGTGT CATCCGAGTG	1380
GCCAGCGGCT ACCTACTGAT GCTTGCCAT GCCTGTTTAA CCATGCTGCG CTGGGACTGC	1440
TCCAAGTCCC AGGGTGCCGT GGGGCTGGCT GCGCTCCTGT TGGTTGCGCT GTCAGTGGCT	1500
GCAGGATTGG GCCTCTGCTC CTTGATTGGC ATTTCTTTTA ATGCTGCGAC AACTCAGGTT	1560
TTGCCGTTTC TTGCTCTTGG TGTTGGTGTG GATGATGTCT TCCTCCTGGC CCATGCATTC	1620
AGTGAAACAG GACAGAATAA GAGGATTCCA TTTGAGGACA GGACTGGGGA GTGCCTCAAG	1680

CGCACCGGAG CCAGCGTGGC CCTCACCTCC ATCAGCAATG TCACCGCCTT CTTCATGGCC	1740
GCATTGATCC CTATCCCTGC CCTGCGAGCG TTCTCCCTCC AGGCTGCTGT GGTGGTGGTA	1800
TTCAATTTTG CTATGGTTCT GCTCATTTTT CCTGCAATTC TCAGCATGGA TTTATACAGA	1860
CGTGAGGACA GAAGATTGGA TATTTTCTGC TGTTTCACAA GCCCCTGTGT CAGCAGGGTG	1920
ATTCAAGTTG AGCCACAGGC CTACACAGAG CCTCACAGTA ACACCCGGTA CAGCCCCCA	1980
CCCCCATA CAAGCCACAG CTTGCCCCAC GAAACCCATA TCACTATGCA GTCCACCGTT	2040
CAGCTCCGCA CAGAGTATGA CCCTCACAG CACGTGTACT ACACCACCGC CGAGCCACGC	2100
TCTGAGATCT CTGTACAGCC TGTTACCGTC ACCCAGGACA ACCTCAGCTG TCAGAGTCCC	2160
GAGAGCACCA GCTCTACCAG GGACCTGCTC TCCAGTTCT CAGACTCCAG CCTCCACTGC	2220
CTCGAGCCCC CCTGCACCAA GTGGACACTC TCTTCGTTTG CAGAGAAGCA CTATGCTCCT	2280
TTCTCCTGA AACCCAAAGC CAAGGTTGTG GTAATCCTTC TTTTCCTGGG CTGCTGGGG	2340
GTCAGCCTTT ATGGGACCAC CCGAGTGAGA GACGGGCTGG ACCTCACGGA CATTGTTCCC	2400
CGGGAAACCA GAGAATATGA CTTCATAGCT GCCCAGTTCA AGTACTTCTC TTTCTACAAC	2460
ATGTATATAG TCACCCAGAA AGCAGACTAC CCGAATATCC AGCACCTACT TTACGACCTT	2520
CATAAGAGTT TCAGCAATGT GAAGTATGTC ATGCTGGAGG AGAACAAGCA ACTTCCCCAA	2580
ATGTGGCTGC ACTACTTTAG AGACTGGCTT CAAGGACTTC AGGATGCATT TGACAGTGAC	2640
TGGGAAACTG GGAGGATCAT GCCAAACAAT TATAAAAATG GATCAGATGA CGGGGTCCTC	2700
GCTTACAAAC TCCTGGTGCA GACTGGCAGC CGAGACAAGC CCATCGACAT TAGTCAGTTG	2760
ACTAAACAGC GTCTGGTAGA CGCAGATGGC ATCATTAATC CGAGCGCTTT CTACATCTAC	2820
CTGACCGCTT GGGTCAGCAA CGACCCTGTA GCTTACGCTG CCTCCCAGGC CAACATCCGG	2880
CCTCACCGGC CGGAGTGGGT CCATGACAAA GCCGACTACA TGCCAGAGAC CAGGCTGAGA	2940
ATCCCAGCAG CAGAGCCCAT CGAGTACGCT CAGTTCCCTT TCTACCTCAA CGGCCTACGA	3000
GACACCTCAG ACTTTGTGGA AGCCATAGAA AAAGTGAGAG TCATCTGTAA CAACTATACG	3060
AGCCTGGGAC TGTCCAGCTA CCCCAATGGC TACCCCTTCC TGTTCTGGGA GCAATACATC	3120
AGCCTGCGCC ACTGGCTGCT GCTATCCATC AGCGTGGTGC TGGCCTGCAC GTTTCTAGTG	3180
TGCGCAGTCT TCCTCCTGAA CCCCTGGACG GCCGGGATCA TTGTCATGGT CCTGGCTCTG	3240
ATGACCGTTG AGCTCTTTGG CATGATGGGC CTCATTGGGA TCAAGCTGAG TGCTGTGCCT	3300
GTGGTCATCC TGATTGCATC TGTTGGCATC GGAGTGGAGT TCACCGTCCA CGTGGCTTTG	3360
GCCTTTCTGA CAGCCATTGG GGACAAGAAC CACAGGGCTA TGCTCGCTCT GGAACACATG	3420
TTTGCTCCCG TTCTGGACGG TGCTGTGTCC ACTCTGCTGG GTGTACTGAT GCTTGACAGG	3480
TCCGAATTTG ATTTTATTGT CAGATACTTC TTTGCCGTCC TGGCCATTCT CACCGTCTTG	3540
GGGGTTCTCA ATGGACTGGT TCTGCTGCCT GTCCTCTTAT CCTTCTTTGG ACCGTGTCCT	3600

GAGGTGTCTC CAGCCAATGG CCTAAACCGA CTGCCCCTC CTTGCGCTGA GCCGCCTCCA 3660  
 AGTGTCTGTC GGTGTGCCGT GCCTCCTGGT CACACGAACA ATGGGTCTGA TTCCTCCGAC 3720  
 TCGGAGTACA GCTCTCAGAC CACGGTGTCT GGCATCAGTG AGGAGCTCAG GCAATACGAA 3780  
 GCACAGCAGG GTGCCGGAGG CCCTGCCCCAC CAAGTGATTG TGAAGCCAC AGAAAACCTT 3840  
 GTCTTTGCCC GGTCCACTGT GGTCCATCCG GACTCCAGAC ATCAGCCTCC CTTGACCCCT 3900  
 CGGCAACAGC CCCACCTGGA CTCTGGCTCC TTGTCCCCTG GACGGCAAGG CCAGCAGCCT 3960  
 CGAAGGGATC CCCCTAGAGA AGGCTTGCGG CCACCCCCCT ACAGACCGCG CAGAGACGCT 4020  
 TTTGAAATTT CTAAGAAG GCATTCTGGC CCTAGCAATA GGGACCGCTC AGGGCCCCGT 4080  
 GGGGCCCGTT CTCACAACCC TCGGAACCCA ACGTCCACCG CCATGGGCAG CTCTGTGCCC 4140  
 AGCTACTGCC AGCCCATCAC CACTGTGACG GCTTCTGCTT CGGTGACTGT TGCTGTGCAT 4200  
 CCCCCGCTG GACCTGGGCG CAACCCCCGA GGGGGGCCCT GTCCAGGCTA TGAGAGCTAC 4260  
 CCTGAGACTG ATCAGGGGT ATTTGAGGAT CCTCATGTGC CTTTTCATGT CAGGTGTGAG 4320  
 AGGAGGGACT CAAAGGTGGA GGTCATAGAG CTACAGGACG TGAATGTGA GGAGAGGCCG 4380  
 TGGGGGAGCA GCTCCAACTG AGGGTAATTA AAATCTGAAG CAAAGAGGCC AAAGATTGGA 4440  
 AAGCCCCGCC CCCACCTCTT TCCAGAACTG CTTGAAGAGA ACTGCTTGGA ATTATGGGAA 4500  
 GGCAGTTCAT TGTTACTGTA ACTGATTGTA TTATTKKGTG AAATATTTCT ATAAATATTT 4560  
 AARAGGTGTA CACATGTAAT ATACATGGAA ATGCTGTACA GTCTATTTCC TGGGGCCTCT 4620  
 CCACTCCTGC CCCAGAGTGG GGAGACCACA GGGGCCCTTT CCCCTGTGTA CATTGGTCTC 4680  
 TGTGCCACAA CCAAGCTTAA CTTAGTTTTA AAAAAATCT CCCAGCATAT GTCGCTGCTG 4740  
 CTTAAATATT GTATAATTTA CTTGTATAAT TCTATGCAAA TATTGCTTAT GTAATAGGAT 4800  
 TATTTGTAAA GGTTCCTGTT TAAAATATTT TAAATTTGCA TATCACAACC CTGTGGTAGG 4860  
 ATGAATTGTT ACTGTTAACT TTTGAACACG CTATGCGTGG TAATTGTTTA ACGAGCAGAC 4920  
 ATGAAGAAAA CAGGTTAATC CCAGTGGCTT CTCTAGGGGT AGTTGTATAT GGTTCGCATG 4980  
 GGTGGATGTG TGTGTGCATG TGACTTTCCA ATGTACTGTA TTGTGGTTTG TTGTTGTTGT 5040  
 TGCTGTTGTT GTTCATTTTG GTGTTTTTGG TTGCTTTGTA TGATCTTAGC TCTGGCCTAG 5100  
 GTGGGCTGGG AAGGTCCAGG TCTTTTTCTG TCGTGATGCT GGTGGAAAGG TGACCCCAAT 5160  
 CATCTGTCCT ATTCTCTGGG ACTATTC 5187

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1311 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met	Val	Ala	Pro	Asp	Ser	Glu	Ala	Pro	Ser	Asn	Pro	Arg	Ile	Thr	Ala	1	5	10	15
Ala	His	Glu	Ser	Pro	Cys	Ala	Thr	Glu	Ala	Arg	His	Ser	Ala	Asp	Leu	20	25	30	
Tyr	Ile	Arg	Thr	Ser	Trp	Val	Asp	Ala	Ala	Leu	Ala	Leu	Ser	Glu	Leu	35	40	45	
Glu	Lys	Gly	Asn	Ile	Glu	Gly	Gly	Arg	Thr	Ser	Leu	Trp	Ile	Arg	Ala	50	55	60	
Trp	Leu	Gln	Glu	Gln	Leu	Phe	Ile	Leu	Gly	Cys	Phe	Leu	Gln	Gly	Asp	65	70	75	80
Ala	Gly	Lys	Val	Leu	Phe	Val	Ala	Ile	Leu	Val	Leu	Ser	Thr	Phe	Cys	85	90	95	
Val	Gly	Leu	Lys	Ser	Ala	Gln	Ile	His	Thr	Arg	Val	Asp	Gln	Leu	Trp	100	105	110	
Val	Gln	Glu	Gly	Gly	Arg	Leu	Glu	Ala	Glu	Leu	Lys	Tyr	Thr	Ala	Gln	115	120	125	
Ala	Leu	Gly	Glu	Ala	Asp	Ser	Ser	Thr	His	Gln	Leu	Val	Ile	Gln	Thr	130	135	140	
Ala	Lys	Asp	Pro	Asp	Val	Ser	Leu	Leu	His	Pro	Gly	Ala	Leu	Leu	Glu	145	150	155	160
His	Leu	Lys	Val	Val	His	Ala	Ala	Thr	Arg	Val	Thr	Val	His	Met	Tyr	165	170	175	
Asp	Ile	Glu	Trp	Arg	Leu	Lys	Asp	Leu	Cys	Tyr	Ser	Pro	Ser	Ile	Pro	180	185	190	
Asp	Phe	Glu	Gly	Tyr	His	His	Ile	Glu	Ser	Ile	Ile	Asp	Asn	Val	Ile	195	200	205	
Pro	Cys	Ala	Ile	Ile	Thr	Pro	Leu	Asp	Cys	Phe	Trp	Glu	Gly	Ser	Lys	210	215	220	
Leu	Leu	Gly	Pro	Asp	Tyr	Pro	Ile	Tyr	Val	Pro	His	Leu	Lys	His	Lys	225	230	235	240
Leu	Gln	Trp	Thr	His	Leu	Asn	Pro	Leu	Glu	Val	Val	Glu	Glu	Val	Lys	245	250	255	
Lys	Leu	Lys	Phe	Gln	Phe	Pro	Leu	Ser	Thr	Ile	Glu	Ala	Tyr	Met	Lys	260	265	270	
Arg	Ala	Gly	Ile	Thr	Ser	Ala	Tyr	Met	Lys	Lys	Pro	Cys	Leu	Asp	Pro	275	280	285	
Thr	Asp	Pro	His	Cys	Pro	Ala	Thr	Ala	Pro	Asn	Lys	Lys	Ser	Gly	His	290	295	300	

16250-07993

Ile Pro Asp Val Ala Ala Glu Leu Ser His Gly Cys Tyr Gly Phe Ala  
 305 310 315 320  
 Ala Ala Tyr Met His Trp Pro Glu Gln Leu Ile Val Gly Gly Ala Thr  
 325 330 335  
 Arg Asn Ser Thr Ser Ala Leu Arg Lys Ala Arg Xaa Leu Gln Thr Val  
 340 345 350  
 Val Gln Leu Met Gly Glu Arg Glu Met Tyr Glu Tyr Trp Ala Asp His  
 355 360 365  
 Tyr Lys Val His Gln Ile Gly Trp Asn Gln Glu Lys Ala Ala Ala Val  
 370 375 380  
 Leu Asp Ala Trp Gln Arg Lys Phe Ala Ala Glu Val Arg Lys Ile Thr  
 385 390 395 400  
 Thr Ser Gly Ser Val Ser Ser Ala Tyr Ser Phe Tyr Pro Phe Ser Thr  
 405 410 415  
 Ser Thr Leu Asn Asp Ile Leu Gly Lys Phe Ser Glu Val Ser Leu Lys  
 420 425 430  
 Asn Ile Ile Leu Gly Tyr Met Phe Met Leu Ile Tyr Val Ala Val Thr  
 435 440 445  
 Leu Ile Gln Trp Arg Asp Pro Ile Arg Ser Gln Ala Gly Val Gly Ile  
 450 455 460  
 Ala Gly Val Leu Leu Leu Ser Ile Thr Val Ala Ala Gly Leu Gly Phe  
 465 470 475 480  
 Cys Ala Leu Leu Gly Ile Pro Phe Asn Ala Ser Ser Thr Gln Ile Val  
 485 490 495  
 Pro Phe Leu Ala Leu Gly Leu Gly Val Gln Asp Met Phe Leu Leu Thr  
 500 505 510  
 His Thr Tyr Val Glu Gln Ala Gly Asp Val Pro Arg Glu Glu Arg Thr  
 515 520 525  
 Gly Leu Val Leu Lys Lys Ser Gly Leu Ser Val Leu Leu Ala Ser Leu  
 530 535 540  
 Cys Asn Val Met Ala Phe Leu Ala Ala Ala Leu Leu Pro Ile Pro Ala  
 545 550 555 560  
 Phe Arg Val Phe Cys Leu Gln Ala Ala Ile Leu Leu Leu Phe Asn Leu  
 565 570 575  
 Gly Ser Ile Leu Leu Val Phe Pro Ala Met Ile Ser Leu Asp Leu Arg  
 580 585 590  
 Arg Arg Ser Ala Ala Arg Ala Asp Leu Leu Cys Cys Leu Met Pro Glu  
 595 600 605  
 Ser Pro Leu Pro Lys Lys Lys Ile Pro Glu Arg Ala Lys Thr Arg Lys  
 610 615 620  
 Asn Asp Lys Thr His Arg Ile Asp Thr Thr Arg Gln Pro Leu Asp Pro  
 625 630 635 640

0016140 002199

Asp	Val	Ser	Glu	Asn	Val	Thr	Lys	Thr	Cys	Cys	Leu	Ser	Val	Ser	Leu
				645					650						655
Thr	Lys	Trp	Ala	Lys	Asn	Gln	Tyr	Ala	Pro	Phe	Ile	Met	Arg	Pro	Ala
			660					665					670		
Val	Lys	Val	Thr	Ser	Met	Leu	Ala	Leu	Ile	Ala	Val	Ile	Leu	Thr	Ser
		675					680					685			
Val	Trp	Gly	Ala	Thr	Lys	Val	Lys	Asp	Gly	Leu	Asp	Leu	Thr	Asp	Ile
	690					695					700				
Val	Pro	Glu	Asn	Thr	Asp	Glu	His	Glu	Phe	Leu	Ser	Arg	Gln	Glu	Lys
705					710					715					720
Tyr	Phe	Gly	Phe	Tyr	Asn	Met	Tyr	Ala	Val	Thr	Gln	Gly	Asn	Phe	Glu
				725					730					735	
Tyr	Pro	Thr	Asn	Gln	Lys	Leu	Leu	Tyr	Glu	Tyr	His	Asp	Gln	Phe	Val
			740					745					750		
Arg	Ile	Pro	Asn	Ile	Ile	Lys	Asn	Asp	Asn	Gly	Gly	Leu	Thr	Lys	Phe
		755					760					765			
Trp	Leu	Ser	Leu	Phe	Arg	Asp	Trp	Leu	Leu	Asp	Leu	Gln	Val	Ala	Phe
	770					775					780				
Asp	Lys	Glu	Val	Ala	Ser	Gly	Cys	Ile	Thr	Gln	Glu	Tyr	Trp	Cys	Lys
785					790					795					800
Asn	Ala	Ser	Asp	Glu	Gly	Ile	Leu	Ala	Tyr	Lys	Leu	Met	Val	Gln	Thr
				805					810					815	
Gly	His	Val	Asp	Asn	Pro	Ile	Asp	Lys	Ser	Leu	Ile	Thr	Ala	Gly	His
			820					825					830		
Arg	Leu	Val	Asp	Lys	Asp	Gly	Ile	Ile	Asn	Pro	Lys	Ala	Phe	Tyr	Asn
		835					840					845			
Tyr	Leu	Ser	Ala	Trp	Ala	Thr	Asn	Asp	Ala	Leu	Ala	Tyr	Gly	Ala	Ser
	850					855					860				
Gln	Gly	Asn	Leu	Lys	Pro	Gln	Pro	Gln	Arg	Trp	Ile	His	Ser	Pro	Glu
865					870					875					880
Asp	Val	His	Leu	Glu	Ile	Lys	Lys	Ser	Ser	Pro	Leu	Ile	Tyr	Thr	Gln
				885					890					895	
Leu	Pro	Phe	Tyr	Leu	Ser	Gly	Leu	Ser	Asp	Thr	Xaa	Ser	Ile	Lys	Thr
			900					905					910		
Leu	Ile	Arg	Ser	Val	Arg	Asp	Leu	Cys	Leu	Lys	Tyr	Glu	Ala	Lys	Gly
		915					920					925			
Leu	Pro	Asn	Phe	Pro	Ser	Gly	Ile	Pro	Phe	Leu	Phe	Trp	Glu	Gln	Tyr
	930					935					940				
Leu	Tyr	Leu	Arg	Thr	Ser	Leu	Leu	Leu	Ala	Leu	Ala	Cys	Ala	Leu	Ala
945					950					955					960
Ala	Val	Phe	Ile	Ala	Val	Met	Val	Leu	Leu	Leu	Asn	Ala	Trp	Ala	Ala
				965						970				975	

Val	Leu	Val	Thr	Leu	Ala	Leu	Ala	Thr	Leu	Val	Leu	Gln	Leu	Gly	
			980					985					990		
Val	Met	Ala	Leu	Leu	Gly	Val	Lys	Leu	Ser	Ala	Met	Pro	Ala	Val	Leu
		995					1000					1005			
Leu	Val	Leu	Ala	Ile	Gly	Arg	Gly	Val	His	Phe	Thr	Val	His	Leu	Cys
	1010					1015					1020				
Leu	Gly	Phe	Val	Thr	Ser	Ile	Gly	Cys	Lys	Arg	Arg	Arg	Ala	Ser	Leu
1025					1030					1035					1040
Ala	Leu	Glu	Ser	Val	Leu	Ala	Pro	Val	Val	His	Gly	Ala	Leu	Ala	Ala
				1045					1050					1055	
Ala	Leu	Ala	Ala	Ser	Met	Leu	Ala	Ala	Ser	Glu	Cys	Gly	Phe	Val	Ala
			1060					1065					1070		
Arg	Leu	Phe	Leu	Arg	Leu	Leu	Leu	Asp	Ile	Val	Phe	Leu	Gly	Leu	Ile
	1075						1080					1085			
Asp	Gly	Leu	Leu	Phe	Phe	Pro	Ile	Val	Leu	Ser	Ile	Leu	Gly	Pro	Ala
	1090					1095					1100				
Ala	Glu	Val	Arg	Pro	Ile	Glu	His	Pro	Glu	Arg	Leu	Ser	Thr	Pro	Ser
1105					1110					1115					1120
Pro	Lys	Cys	Ser	Pro	Ile	His	Pro	Arg	Lys	Ser	Ser	Ser	Ser	Ser	Gly
				1125					1130						1135
Gly	Gly	Asp	Lys	Ser	Ser	Arg	Thr	Ser	Lys	Ser	Ala	Pro	Arg	Pro	Cys
			1140					1145					1150		
Ala	Pro	Ser	Leu	Thr	Thr	Ile	Thr	Glu	Glu	Pro	Ser	Ser	Trp	His	Ser
		1155					1160					1165			
Ser	Ala	His	Ser	Val	Gln	Ser	Ser	Met	Gln	Ser	Ile	Val	Val	Gln	Pro
	1170					1175					1180				
Glu	Val	Val	Val	Glu	Thr	Thr	Thr	Tyr	Asn	Gly	Ser	Asp	Ser	Ala	Ser
1185					1190					1195					1200
Gly	Arg	Ser	Thr	Pro	Thr	Lys	Ser	Ser	His	Gly	Gly	Ala	Ile	Thr	Thr
				1205					1210					1215	
Thr	Lys	Val	Thr	Ala	Thr	Ala	Asn	Ile	Lys	Val	Glu	Val	Val	Thr	Pro
			1220					1225					1230		
Ser	Asp	Arg	Lys	Ser	Arg	Arg	Ser	Tyr	His	Tyr	Tyr	Asp	Arg	Arg	Arg
	1235						1240					1245			
Asp	Arg	Asp	Glu	Asp	Arg	Asp	Arg	Asp	Arg	Glu	Arg	Asp	Arg	Asp	Arg
	1250					1255					1260				
Asp	Arg	Asp	Arg	Asp	Arg	Asp	Arg	Asp	Arg	Asp	Arg	Asp	Arg	Asp	Arg
1265					1270					1275					1280
Glu	Arg	Ser	Arg	Glu	Arg	Asp	Arg	Arg	Asp	Arg	Tyr	Arg	Asp	Glu	Arg
				1285					1290					1295	
Asp	His	Arg	Ala	Ser	Pro	Arg	Glu	Lys	Arg	Gln	Arg	Phe	Trp	Thr	
			1300					1305							

## (2) INFORMATION FOR SEQ ID NO:5:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 4434 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CGAAACAAGA GAGCGAGTGA GAGTAGGGAG AGCGTCTGTG TTGTGTGTTG AGTGTGCCCC	60
ACGCACACAG GCGCAAAACA GTGCACACAG ACGCCCGCTG GGCAAGAGAG AGTGAGAGAG	120
AGAAACAGCG GCGCGCGCTC GCCTAATGAA GTTGTGTTGCC TGGCTGGCGT GCCGCATCCA	180
CGAGATACAG ATACATCTCT CATGGACCGC GACAGCCTCC CACGCGTTCC GGACACACAC	240
GGCGATGTGG TCGATGAGAA ATTATTCTCG GATCTTTACA TACGCACCAG CTGGGTGGAC	300
GCGCAAGTGG CGCTCGATCA GATAGATAAG GGCAAAGCGC GTGGCAGCCG CACGGCGATC	360
TATCTGCGAT CAGTATTCCA GTCCACCTC GAAACCCCTG GCAGCTCCGT GCAAAAGCAC	420
GCGGGCAAGG TGCTATTCGT GGCTATCCTG GTGCTGAGCA CCTTCTGCGT CGGCCTGAAG	480
AGCGCCCAGA TCCACTCCAA GGTGCACCAG CTGTGGATCC AGGAGGGCGG CCGGCTGGAG	540
GCGGAACTGG CCTACACACA GAAGACGATC GGCGAGGACG AGTCGGCCAC GCATCAGCTG	600
CTCATTCAGA CGACCCACGA CCCGAACGCC TCCGTCCTGC ATCCGCAGGC GCTGCTTGCC	660
CACCTGGAGG TCCTGGTCAA GGCCACCGCC GTCAAGGTGC ACCTCTACGA CACCGAATGG	720
GGGCTGCGCG ACATGTGCAA CATGCCGAGC ACGCCCTCCT TCGAGGGCAT CTACTACATC	780
GAGCAGATCC TGCGCCACCT CATTCCGTGC TCGATCATCA CGCCGCTGGA CTGTTTCTGG	840
GAGGGAAGCC AGCTGTTGGG TCCGGAATCA GCGGTCGTTA TACCAGGCCT CAACCAACGA	900
CTCCTGTGGA CCACCCTGAA TCCCGCCTCT GTGATGCAGT ATATGAAACA AAAGATGTCC	960
GAGGAAAAGA TCAGCTTCGA CTTGAGACC GTGGAGCAGT ACATGAAGCG TCGCGCCATT	1020
GGCAGTGGCT ACATGGAGAA GCCCTGCCTG AACCCTACTGA ATCCCAATTG CCCGGACACG	1080
GCACCGAACA AGAACAGCAC CCAGCCGCCG GATGTGGGAG CCATCCTGTC CGGAGGCTGC	1140
TACGGTTATG CCGCGAAGCA CATGCACTGG CCGGAGGAGC TGATTGTGGG CGGACGGAAG	1200
AGGAACCGCA GCGGACACTT GAGGAAGGCC CAGGCCCTGC AGTCGGTGGT GCAGCTGATG	1260
ACCGAGAAGG AAATGTACGA CCAGTGGCAG GACAACTACA AGGTGCACCA TCTTGATGG	1320
ACGCAGGAGA AGGCAGCGGA GGTTTTGAAC GCCTGGCAGC GCAACTTTTC GCGGGAGGTG	1380
GAACAGCTGC TACGTAAACA GTCGAGAATT GCCACCAACT ACGATATCTA CGTGTTACAGC	1440

TCGGCTGCAC TGGATGACAT CCTGGCCAAG TTCTCCCATC CCAGCGCCTT GTCCATTGTC 1500  
 ATCGGCGTGG CCGTCACCGT TTTGTATGCC TTTTGACACG TCCTCCGCTG GAGGGACCCC 1560  
 GTCCGTGGCC AGAGCAGTGT GGGCGTGGCC GGAGTTCTGC TCATGTGCTT CAGTACCGCC 1620  
 GCCGGATTGG GATTGTCAGC CCTGCTCGGT ATCGTTTTCA ATGCGCTGAC CGCTGCCTAT 1680  
 GCGGAGAGCA ATCGGCGGGA GCAGACCAAG CTGATTCTCA AGAACGCCAG CACCCAGGTG 1740  
 GTTCCGTTTT TGGCCCTTGG TCTGGGCGTC GATCACATCT TCATAGTGGG ACCGAGCATC 1800  
 CTGTTCAAGT CCTGCAGCAC CGCAGGATCC TTCTTTGCGG CCGCCTTTAT TCCGGTGCCG 1860  
 GCTTTGAAGG TATTCTGTCT GCAGGCTGCC ATCGTAATGT GCTCCAATTT GGCAGCGGCT 1920  
 CTATTGGTTT TTCCGGCCAT GATTTTCGTT GATCTACGGA GACGTACCGC CGGCAGGGCG 1980  
 GACATCTTCT GCTGCTGTTT TCCGGTGTGG AAGGAACAGC CGAAGGTGGC ACCTCCGGTG 2040  
 CTGCCGCTGA ACAACAACAA CGGGCGCGGG GCCCGGCATC CGAAGAGCTG CAACAACAAC 2100  
 AGGGTGCCGC TGCCCGCCCA GAATCCTCTG CTGGAACAGA GGGCAGACAT CCCTGGGAGC 2160  
 AGTCACTCAC TGGCGTCCTT CTCCCTGGCA ACCTTCGCCT TTCAGACTA CACTCCCTTC 2220  
 CTCATGCGCA GCTGGGTGAA GTTCCTGACC GTTATGGGTT TCCTGGCGGC CCTCATATCC 2280  
 AGCTTGATG CCTCCACGCG CCTTCAGGAT GGCCTGGACA TTATTGATCT GGTGCCCCAAG 2340  
 GACAGCAACG AGCACAAGTT CCTGGATGCT CAAACTCGGC TCTTTGGCTT CTACAGCATG 2400  
 TATGCGGTTA CCCAGGGCAA CTTTGAATAT CCCACCCAGC AGCAGTTGCT CAGGGACTAC 2460  
 CATGATTCCCT TTGTGCGGGT GCCACATGTG ATCAAGAATG ATAACGGTGG ACTGCCGGAC 2520  
 TTCTGGCTGC TGCTCTTCAG CGAGTGCGTG GGTAATCTGC AAAAGATATT CGACGAGGAA 2580  
 TACCGCGACG GACGGCTGAC CAAGGAGTGC TGGTTCCCAA ACGCCAGCAG CGATGCCATC 2640  
 CTGGCCTACA AGCTAATCGT GCAAACCGGC CATGTGGACA ACCCCGTGGA CAAGGAACTG 2700  
 GTGCTACCA ATCGCCTGGT CAACAGCGAT GGCATCATCA ACCAACGCGC CTTCTACAAC 2760  
 TATCTGTGCG CATGGGCCAC CAACGACGTC TTCGCCTACG GAGCTTCTCA GGGCAAATTG 2820  
 TATCCGGAAC CGCGCCAGTA TTTTCACCAA CCCAACGAGT ACGATCTTAA GATACCCAAG 2880  
 AGTCTGCCAT TGGTCTACGC TCAGATGCCC TTTTACCTCC ACGGACTAAC AGATACCTCG 2940  
 CAGATCAAGA CCCTGATAGG TCATATTCGC GACCTGAGCG TCAAGTACGA GGGCTTCGGC 3000  
 CTGCCCAACT ATCCATCGGG CATTCCCTTC ATCTTCTGGG AGCAGTACAT GACCCTGCGC 3060  
 TCCTCACTGG CCATGATCCT GGCCTGCGTG CTACTCGCCG CCCTGGTGCT GGTCTCCCTG 3120  
 CTCCTGCTCT CCGTTTGGGC CGCCGTTCTC GTGATCCTCA GCGTTCTGGC CTCGCTGGCC 3180  
 CAGATCTTTG GGGCCATGAC TCTGCTGGGC ATCAAATCTC CGGCCATTCC GGCAGTCATA 3240  
 CTCATCCTCA GCGTGGGCAT GATGCTGTGC TTCAATGTGC TGATATCACT GGGCTTCATG 3300  
 ACATCCGTTG GCAACCGACA GCGCCGCGTC CAGCTGAGCA TGCAGATGTC CCTGGGACCA 3360

CTTGTCACG GCATGCTGAC CTCCGGAGTG GCCGTGTTCA TGCTCTCCAC GTCGCCCTTT 3420  
 GAGTTTGTGA TCCGGCACTT CTGCTGGCTT CTGCTGGTGG TCTTATGCGT TGGCGCCTGC 3480  
 AACAGCCTTT TGGTGTTCCT CATCCTACTG AGCATGGTGG GACCGGAGGC GGAGCTGGTG 3540  
 CCGCTGGAGC ATCCAGACCG CATATCCACG CCCTCTCCGC TGCCCGTGCG CAGCAGCAAG 3600  
 AGATCGGGCA AATCCTATGT GGTGCAGGGA TCGCGATCCT CGCGAGGCAG CTGCCAGAAG 3660  
 TCGCATCACC ACCACCACAA AGACCTTAAT GATCCATCGC TGACGACGAT CACCGAGGAG 3720  
 CCGCAGTCGT GGAAGTCCAG CAACTCGTCC ATCCAGATGC CCAATGATTG GACCTACCAG 3780  
 CCGCGGGAAC AGCGACCCGC CTCCTACGCG GCCCGCCCCC CCGCCTATCA CAAGGCCGCC 3840  
 GCCCAGCAGC ACCACCAGCA TCAGGGCCCG CCCACAACGC CCCC GCCTCC CTTCCCGACG 3900  
 GCCTATCCGC CGGAGCTGCA GAGCATCGTG GTGCAGCCGG AGGTGACGGT GGAGACGACG 3960  
 CACTCGGACA GCAACACCAC CAAGGTGACG GCCACGGCCA ACATCAAGGT GGAGCTGGCC 4020  
 ATGCCCCGCA GGGCGGTGCG CAGCTATAAC TTTACGAGTT AGCACTAGCA CTAGTTCCTG 4080  
 TAGCTATTAG GACGTATCTT TAGACTCTAG CCTAAGCCGT AACCCCTATTT GTATCTGTAA 4140  
 AATCGATTTG TCCAGCGGGT CTGCTGAGGA TTTCGTTCTC ATGGATTCTC ATGGATTCTC 4200  
 ATGGATGCTT AAATGGCATG GTAATTGGCA AAATATCAAT TTTTGTGTCT CAAAAGATG 4260  
 CATTAGCTTA TGGTTTCAAG ATACATTTTT AAAGAGTCCG CCAGATATTT ATATAAAAAA 4320  
 AATCCAAAAT CGACGTATCC ATGAAAATTG AAAAGCTAAG CAGACCCGTA TGTATGTATA 4380  
 TGTGTATGCA TGTTAGTTAA TTTCCGAAG TCCGGTATTT ATAGCAGCTG CCTT 4434

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1285 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met	Asp	Arg	Asp	Ser	Leu	Pro	Arg	Val	Pro	Asp	Thr	His	Gly	Asp	Val
1				5					10					15	
Val	Asp	Glu	Lys	Leu	Phe	Ser	Asp	Leu	Tyr	Ile	Arg	Thr	Ser	Trp	Val
			20					25					30		
Asp	Ala	Gln	Val	Ala	Leu	Asp	Gln	Ile	Asp	Lys	Gly	Lys	Ala	Arg	Gly
		35					40					45			
Ser	Arg	Thr	Ala	Ile	Tyr	Leu	Arg	Ser	Val	Phe	Gln	Ser	His	Leu	Glu
	50					55					60				

Thr	Leu	Gly	Ser	Ser	Val	Gln	Lys	His	Ala	Gly	Lys	Val	Leu	Phe	Val
65					70					75					80
Ala	Ile	Leu	Val	Leu	Ser	Thr	Phe	Cys	Val	Gly	Leu	Lys	Ser	Ala	Gln
				85					90					95	
Ile	His	Ser	Lys	Val	His	Gln	Leu	Trp	Ile	Gln	Glu	Gly	Gly	Arg	Leu
			100					105					110		
Glu	Ala	Glu	Leu	Ala	Tyr	Thr	Gln	Lys	Thr	Ile	Gly	Glu	Asp	Glu	Ser
		115					120					125			
Ala	Thr	His	Gln	Leu	Leu	Ile	Gln	Thr	Thr	His	Asp	Pro	Asn	Ala	Ser
	130					135					140				
Val	Leu	His	Pro	Gln	Ala	Leu	Leu	Ala	His	Leu	Glu	Val	Leu	Val	Lys
145					150					155					160
Ala	Thr	Ala	Val	Lys	Val	His	Leu	Tyr	Asp	Thr	Glu	Trp	Gly	Leu	Arg
				165					170					175	
Asp	Met	Cys	Asn	Met	Pro	Ser	Thr	Pro	Ser	Phe	Glu	Gly	Ile	Tyr	Tyr
			180					185					190		
Ile	Glu	Gln	Ile	Leu	Arg	His	Leu	Ile	Pro	Cys	Ser	Ile	Ile	Thr	Pro
		195					200					205			
Leu	Asp	Cys	Phe	Trp	Glu	Gly	Ser	Gln	Leu	Leu	Gly	Pro	Glu	Ser	Ala
	210					215					220				
Val	Val	Ile	Pro	Gly	Leu	Asn	Gln	Arg	Leu	Leu	Trp	Thr	Thr	Leu	Asn
225					230					235					240
Pro	Ala	Ser	Val	Met	Gln	Tyr	Met	Lys	Gln	Lys	Met	Ser	Glu	Glu	Lys
				245					250					255	
Ile	Ser	Phe	Asp	Phe	Glu	Thr	Val	Glu	Gln	Tyr	Met	Lys	Arg	Ala	Ala
			260					265					270		
Ile	Gly	Ser	Gly	Tyr	Met	Glu	Lys	Pro	Cys	Leu	Asn	Pro	Leu	Asn	Pro
		275					280					285			
Asn	Cys	Pro	Asp	Thr	Ala	Pro	Asn	Lys	Asn	Ser	Thr	Gln	Pro	Pro	Asp
	290					295					300				
Val	Gly	Ala	Ile	Leu	Ser	Gly	Gly	Cys	Tyr	Gly	Tyr	Ala	Ala	Lys	His
305					310					315					320
Met	His	Trp	Pro	Glu	Glu	Leu	Ile	Val	Gly	Gly	Arg	Lys	Arg	Asn	Arg
				325					330					335	
Ser	Gly	His	Leu	Arg	Lys	Ala	Gln	Ala	Leu	Gln	Ser	Val	Val	Gln	Leu
			340					345					350		
Met	Thr	Glu	Lys	Glu	Met	Tyr	Asp	Gln	Trp	Gln	Asp	Asn	Tyr	Lys	Val
		355					360					365			
His	His	Leu	Gly	Trp	Thr	Gln	Glu	Lys	Ala	Ala	Glu	Val	Leu	Asn	Ala
	370					375					380				
Trp	Gln	Arg	Asn	Phe	Ser	Arg	Glu	Val	Glu	Gln	Leu	Leu	Arg	Lys	Gln
385					390					395					400

Ser Arg Ile Ala Thr Asn Tyr Asp Ile Tyr Val Phe Ser Ser Ala Ala  
 405 410 415  
 Leu Asp Asp Ile Leu Ala Lys Phe Ser His Pro Ser Ala Leu Ser Ile  
 420 425 430  
 Val Ile Gly Val Ala Val Thr Val Leu Tyr Ala Phe Cys Thr Leu Leu  
 435 440 445  
 Arg Trp Arg Asp Pro Val Arg Gly Gln Ser Ser Val Gly Val Ala Gly  
 450 455 460  
 Val Leu Leu Met Cys Phe Ser Thr Ala Ala Gly Leu Gly Leu Ser Ala  
 465 470 475 480  
 Leu Leu Gly Ile Val Phe Asn Ala Leu Thr Ala Ala Tyr Ala Glu Ser  
 485 490 495  
 Asn Arg Arg Glu Gln Thr Lys Leu Ile Leu Lys Asn Ala Ser Thr Gln  
 500 505 510  
 Val Val Pro Phe Leu Ala Leu Gly Leu Gly Val Asp His Ile Phe Ile  
 515 520 525  
 Val Gly Pro Ser Ile Leu Phe Ser Ala Cys Ser Thr Ala Gly Ser Phe  
 530 535 540  
 Phe Ala Ala Ala Phe Ile Pro Val Pro Ala Leu Lys Val Phe Cys Leu  
 545 550 555 560  
 Gln Ala Ala Ile Val Met Cys Ser Asn Leu Ala Ala Ala Leu Leu Val  
 565 570 575  
 Phe Pro Ala Met Ile Ser Leu Asp Leu Arg Arg Arg Thr Ala Gly Arg  
 580 585 590  
 Ala Asp Ile Phe Cys Cys Cys Phe Pro Val Trp Lys Glu Gln Pro Lys  
 595 600 605  
 Val Ala Pro Pro Val Leu Pro Leu Asn Asn Asn Asn Gly Arg Gly Ala  
 610 615 620  
 Arg His Pro Lys Ser Cys Asn Asn Asn Arg Val Pro Leu Pro Ala Gln  
 625 630 635 640  
 Asn Pro Leu Leu Glu Gln Arg Ala Asp Ile Pro Gly Ser Ser His Ser  
 645 650 655  
 Leu Ala Ser Phe Ser Leu Ala Thr Phe Ala Phe Gln His Tyr Thr Pro  
 660 665 670  
 Phe Leu Met Arg Ser Trp Val Lys Phe Leu Thr Val Met Gly Phe Leu  
 675 680 685  
 Ala Ala Leu Ile Ser Ser Leu Tyr Ala Ser Thr Arg Leu Gln Asp Gly  
 690 695 700  
 Leu Asp Ile Ile Asp Leu Val Pro Lys Asp Ser Asn Glu His Lys Phe  
 705 710 715 720  
 Leu Asp Ala Gln Thr Arg Leu Phe Gly Phe Tyr Ser Met Tyr Ala Val  
 725 730 735  
 Thr Gln Gly Asn Phe Glu Tyr Pro Thr Gln Gln Gln Leu Leu Arg Asp

00916140 082197  
 00916140 082197

740										745										750										
Tyr	His	Asp	Ser	Phe	Arg	Val	Pro	His	Val	Ile	Lys	Asn	Asp	Asn	Gly															
		755					760					765																		
Gly	Leu	Pro	Asp	Phe	Trp	Leu	Leu	Leu	Phe	Ser	Glu	Trp	Leu	Gly	Asn															
	770					775					780																			
Leu	Gln	Lys	Ile	Phe	Asp	Glu	Glu	Tyr	Arg	Asp	Gly	Arg	Leu	Thr	Lys															
785					790					795					800															
Glu	Cys	Trp	Phe	Pro	Asn	Ala	Ser	Ser	Asp	Ala	Ile	Leu	Ala	Tyr	Lys															
				805					810					815																
Leu	Ile	Val	Gln	Thr	Gly	His	Val	Asp	Asn	Pro	Val	Asp	Lys	Glu	Leu															
			820					825					830																	
Val	Leu	Thr	Asn	Arg	Leu	Val	Asn	Ser	Asp	Gly	Ile	Ile	Asn	Gln	Arg															
			835				840						845																	
Ala	Phe	Tyr	Asn	Tyr	Leu	Ser	Ala	Trp	Ala	Thr	Asn	Asp	Val	Phe	Ala															
		850				855					860																			
Tyr	Gly	Ala	Ser	Gln	Gly	Lys	Leu	Tyr	Pro	Glu	Pro	Arg	Gln	Tyr	Phe															
865					870					875					880															
His	Gln	Pro	Asn	Glu	Tyr	Asp	Leu	Lys	Ile	Pro	Lys	Ser	Leu	Pro	Leu															
				885					890					895																
Val	Tyr	Ala	Gln	Met	Pro	Phe	Tyr	Leu	His	Gly	Leu	Thr	Asp	Thr	Ser															
			900					905					910																	
Gln	Ile	Lys	Thr	Leu	Ile	Gly	His	Ile	Arg	Asp	Leu	Ser	Val	Lys	Tyr															
		915					920					925																		
Glu	Gly	Phe	Gly	Leu	Pro	Asn	Tyr	Pro	Ser	Gly	Ile	Pro	Phe	Ile	Phe															
	930					935					940																			
Trp	Glu	Gln	Tyr	Met	Thr	Leu	Arg	Ser	Ser	Leu	Ala	Met	Ile	Leu	Ala															
945					950					955					960															
Cys	Val	Leu	Leu	Ala	Ala	Leu	Val	Leu	Val	Ser	Leu	Leu	Leu	Leu	Ser															
				965					970						975															
Val	Trp	Ala	Ala	Val	Leu	Val	Ile	Leu	Ser	Val	Leu	Ala	Ser	Leu	Ala															
			980					985					990																	
Gln	Ile	Phe	Gly	Ala	Met	Thr	Leu	Leu	Gly	Ile	Lys	Leu	Ser	Ala	Ile															
		995					1000					1005																		
Pro	Ala	Val	Ile	Leu	Ile	Leu	Ser	Val	Gly	Met	Met	Leu	Cys	Phe	Asn															
		1010				1015					1020																			
Val	Leu	Ile	Ser	Leu	Gly	Phe	Met	Thr	Ser	Val	Gly	Asn	Arg	Gln	Arg															
1025					1030					1035					1040															
Arg	Val	Gln	Leu	Ser	Met	Gln	Met	Ser	Leu	Gly	Pro	Leu	Val	His	Gly															
				1045					1050					1055																
Met	Leu	Thr	Ser	Gly	Val	Ala	Val	Phe	Met	Leu	Ser	Thr	Ser	Pro	Phe															
			1060					1065					1070																	
Glu	Phe	Val	Ile	Arg	His	Phe	Cys	Trp	Leu	Leu	Leu	Val	Val	Leu	Cys															

00946440-00949

1075	1080	1085
Val Gly Ala Cys Asn Ser Leu Leu Val Phe Pro Ile Leu Leu Ser Met 1090	1095	1100
Val Gly Pro Glu Ala Glu Leu Val Pro Leu Glu His Pro Asp Arg Ile 1105	1110	1115 1120
Ser Thr Pro Ser Pro Leu Pro Val Arg Ser Ser Lys Arg Ser Gly Lys 1125	1130	1135
Ser Tyr Val Val Gln Gly Ser Arg Ser Ser Arg Gly Ser Cys Gln Lys 1140	1145	1150
Ser His His His His His Lys Asp Leu Asn Asp Pro Ser Leu Thr Thr 1155	1160	1165
Ile Thr Glu Glu Pro Gln Ser Trp Lys Ser Ser Asn Ser Ser Ile Gln 1170	1175	1180
Met Pro Asn Asp Trp Thr Tyr Gln Pro Arg Glu Gln Arg Pro Ala Ser 1185	1190	1195 1200
Tyr Ala Ala Pro Pro Pro Ala Tyr His Lys Ala Ala Ala Gln Gln His 1205	1210	1215
His Gln His Gln Gly Pro Pro Thr Thr Pro Pro Pro Pro Phe Pro Thr 1220	1225	1230
Ala Tyr Pro Pro Glu Leu Gln Ser Ile Val Val Gln Pro Glu Val Thr 1235	1240	1245
Val Glu Thr Thr His Ser Asp Ser Asn Thr Thr Lys Val Thr Ala Thr 1250	1255	1260
Ala Asn Ile Lys Val Glu Leu Ala Met Pro Gly Arg Ala Val Arg Ser 1265	1270	1275 1280
Tyr Asn Phe Thr Ser 1285		

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 345 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

AAGGTCCATC AGCTTTGGAT ACAGGAAGGT GGTTCGCTCG AGCATGAGCT AGCCTACACG	60
CAGAAATCGC TCGGCGAGAT GGACTCCTCC ACGCACCAGC TGCTAATCCA AACNCCCAAA	120
GATATGGACG CCTCGATACT GCACCCGAAC GCGCTACTGA CGCACCTGGA CGTGGTGAAG	180
AAAGCGATCT CGGTGACGGT GCACATGTAC GACATCACGT GGAGNCTCAA GGACATGTGC	240

TACTCGCCCA GCATACCGAG NTTGATACG CACTTTATCG AGCAGATCTT CGAGAACATC 300  
 ATACCGTGCG CGATCATCAC GCCGCTGGAT TGCTTTTGGG AGGGA 345

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 115 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Lys Val His Gln Leu Trp Ile Gln Glu Gly Gly Ser Leu Glu His Glu  
 1 5 10 15  
 Leu Ala Tyr Thr Gln Lys Ser Leu Gly Glu Met Asp Ser Ser Thr His  
 20 25 30  
 Gln Leu Leu Ile Gln Thr Pro Lys Asp Met Asp Ala Ser Ile Leu His  
 35 40 45  
 Pro Asn Ala Leu Leu Thr His Leu Asp Val Val Lys Lys Ala Ile Ser  
 50 55 60  
 Val Thr Val His Met Tyr Asp Ile Thr Trp Xaa Leu Lys Asp Met Cys  
 65 70 75 80  
 Tyr Ser Pro Ser Ile Pro Xaa Phe Asp Thr His Phe Ile Glu Gln Ile  
 85 90 95  
 Phe Glu Asn Ile Ile Pro Cys Ala Ile Ile Thr Pro Leu Asp Cys Phe  
 100 105 110  
 Trp Glu Gly  
 115

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 5187 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GGGTCTGTCA CCCGAGCCG GAGTCCCCGG CGGCCAGCAG CGTCCTCGCG AGCCGAGCGC 60  
 CCAGGCGCGC CCGGAGCCCG CGGCGGCGGC GGCAACATGG CCTCGGCTGG TAACGCCGCC 120

09915140.0894

GGGGCCCTGG GCAGGCAGGC CGGCGGCGGG AGGCGCAGAC GGACCGGGGG ACCGCACCGC	180
GCCGCGCCGG ACCGGGACTA TCTGCACCGG CCCAGCTACT GCGACGCCGC CTTGCTCTG	240
GAGCAGATTT CCAAGGGGAA GGCTACTGGC CGGAAAGCGC CGCTGTGGCT GAGAGCGAAG	300
TTTCAGAGAC TCTTATTTAA ACTGGGTTGT TACATTCAAA AGAACTGCGG CAAGTTTTTG	360
GTTGTGGGTC TCCTCATATT TGGGGCCTTC GCTGTGGGAT TAAAGGCAGC TAATCTCGAG	420
ACCAACGTGG AGGAGCTGTG GGTGGAAGTT GGTGGACGAG TGAGTCGAGA ATTAAATTAT	480
ACCCGTCAGA AGATAGGAGA AGAGGCTATG TTTAATCCTC AACTCATGAT ACAGACTCCA	540
AAAGAAGAAG GCGCTAATGT TCTGACCACA GAGGCTCTCC TGCAACACCT GGA CTCAGCA	600
CTCCAGGCCA GTCGTGTGCA CGTCTACATG TATAACAGGC AATGGAAGTT GGAACATTTG	660
TGCTACAAAT CAGGGGAACT TATCACGGAG ACAGGTTACA TGGATCAGAT AATAGAATAC	720
CTTTACCCCTT GCTTAATCAT TACACCTTTG GACTGCTTCT GGGAAGGGGC AAAGCTACAG	780
TCCGGGACAG CATACTCCT AGGTAAGCCT CCTTTACGGT GGACAACTT TGACCCCTTG	840
GAATTCCTAG AAGAGTTAAA GAAAATAAAC TACCAAGTGG ACAGCTGGGA GGAAATGCTG	900
AATAAGCCG AAGTTGGCCA TGGGTACATG GACCGGCCTT GCCTCAACCC AGCCGACCCA	960
GATTGCCCTG CCACAGCCCC TAACAAAAAT TCAACCAAAC CTCTTGATGT GGCCCTTGTT	1020
TTGAATGGTG GATGTCAAGG TTTATCCAGG AAGTATATGC ATTGGCAGGA GGAGTTGATT	1080
GTGGGTGGTA CCGTCAAGAA TGCCACTGGA AAACCTGTCA GCGCTCACGC CCTGCAAACC	1140
ATGTTCCAGT TAATGACTCC CAAGCAAATG TATGAACACT TCAGGGGCTA CGACTATGTC	1200
TCTCACATCA ACTGGAATGA AGACAGGGCA GCCGCCATCC TGGAGGCCTG GCAGAGGACT	1260
TACGTGGAGG TGTTTCATCA AAGTGTCGCC CCAAACCTCCA CTCAAAGGT GCTTCCCTTC	1320
ACAACCACGA CCCTGGACGA CATCCTAAAA TCCTTCTCTG ATGTCAGTGT CATCCGAGTG	1380
GCCAGCGGCT ACCTACTGAT GCTTGCCTAT GCCTGTTTAA CCATGCTGCG CTGGGACTGC	1440
TCCAAGTCCC AGGGTGCCGT GGGGCTGGCT GCGTCTCTGT TGTTGCGCT GTCAGTGGCT	1500
GCAGGATTGG GCCTCTGCTC CTTGATTGGC ATTTCTTTTA ATGCTGCGAC AACTCAGGTT	1560
TTGCCGTTTC TTGCTCTTGG TGTGGTGTG GATGATGTCT TCCTCCTGGC CCATGCATTC	1620
AGTGAAACAG GACAGAATAA GAGGATTCCA TTTGAGGACA GGACTGGGGA GTGCCTCAAG	1680
CGCACC GGAG CCAGCGTGGC CCTCACCCTCC ATCAGCAATG TCACCGCCTT CTTCATGGCC	1740
GCATTGATCC CTATCCCTGC CCTGCGAGCG TTCTCCCTCC AGGCTGCTGT GGTGGTGGTA	1800
TTCAATTTTG CTATGGTTCT GCTCATTTTT CCTGCAATTC TCAGCATGGA TTTATACAGA	1860
CGTGAGGACA GAAGATTGGA TATTTTCTGC TGTTTCACAA GCCCCTGTGT CAGCAGGGTG	1920
ATTCAAGTTG AGCCACAGGC CTACACAGAG CCTCACAGTA ACACCCGGTA CAGCCCCCA	1980
CCCCCATACA CCAGCCACAG CTTGCCCCAC GAAACCCATA TCACTATGCA GTCCACCGTT	2040

CAGCTCCGCA	CAGAGTATGA	CCCTCACACG	CACGTGTACT	ACACCACCGC	CGAGCCACGC	2100
TCTGAGATCT	CTGTACAGCC	TGTTACCGTC	ACCCAGGACA	ACCTCAGCTG	TCAGAGTCCC	2160
GAGAGCACCA	GCTCTACCAG	GGACCTGCTC	TCCCAGTTCT	CAGACTCCAG	CCTCCACTGC	2220
CTCGAGCCCC	CCTGCACCAA	GTGGACACTC	TCTTCGTTTG	CAGAGAAGCA	CTATGCTCCT	2280
TTCTCTCTGA	AACCCAAAGC	CAAGGTTGTG	GTAATCCTTC	TTTTCTCTGGG	CTTGCTGGGG	2340
GTCAGCCTTT	ATGGGACCAC	CCGAGTGAGA	GACGGGCTGG	ACCTCACGGA	CATTGTTCCC	2400
CGGGAAACCA	GAGAATATGA	CTTCATAGCT	GCCCAGTTCA	AGTACTTCTC	TTTCTACAAC	2460
ATGTATATAG	TCACCCAGAA	AGCAGACTAC	CCGAATATCC	AGCACCTACT	TTACGACCTT	2520
CATAAGAGTT	TCAGCAATGT	GAAGTATGTC	ATGCTGGAGG	AGAACAAGCA	ACTTCCCCAA	2580
ATGTGGCTGC	ACTACTTTAG	AGACTGGCTT	CAAGGACTTC	AGGATGCATT	TGACAGTGAC	2640
TGGGAAACTG	GGAGGATCAT	GCCAAACAAT	TATAAAAATG	GATCAGATGA	CGGGGTCTCT	2700
GCTTACAAAC	TCCTGGTGCA	GA CTGGCAGC	CGAGACAAGC	CCATCGACAT	TAGTCAGTTG	2760
ACTAAACAGC	GTCTGGTAGA	CGCAGATGGC	ATCATTAAATC	CGAGCGCTTT	CTACATCTAC	2820
CTGACCGCTT	GGGTCAGCAA	CGACCCTGTA	GCTTACGCTG	CCTCCCAGGC	CAACATCCGG	2880
CCTCACCGGC	CGGAGTGGGT	CCATGACAAA	GCCGACTACA	TGCCAGAGAC	CAGGCTGAGA	2940
ATCCCAGCAG	CAGAGCCCAT	CGAGTACGCT	CAGTTCCCTT	TCTACCTCAA	CGGCCTACGA	3000
GACACCTCAG	ACTTTGTGGA	AGCCATAGAA	AAAGTGAGAG	TCATCTGTAA	CAACTATACG	3060
AGCCTGGGAC	TGTCCAGCTA	CCCCAATGGC	TACCCCTTCC	TGTTCTGGGA	GCAATACATC	3120
AGCCTGCGCC	ACTGGCTGCT	GCTATCCATC	AGCGTGGTGC	TGGCCTGCAC	GTTTCTAGTG	3180
TGCGCAGTCT	TCCTCCTGAA	CCCCTGGACG	GCCGGGATCA	TTGTCATGGT	CCTGGCTCTG	3240
ATGACCGTTG	AGCTCTTTGG	CATGATGGGC	CTCATTGGGA	TCAAAGCTGAG	TGCTGTGCCT	3300
GTGGTCATCC	TGATTGCATC	TGTTGGCATC	GGAGTGGAGT	TCACCGTCCA	CGTGGCTTTG	3360
GCCTTTCTGA	CAGCCATTGG	GGACAAGAAC	CACAGGGCTA	TGCTCGCTCT	GGAACACATG	3420
TTTGCTCCCG	TTCTGGACGG	TGCTGTGTCC	ACTCTGCTGG	GTGTACTGAT	GCTTGCAGGG	3480
TCCGAATTTG	ATTTCAATTGT	CAGATACTTC	TTTGCCGTCC	TGGCCATTCT	CACCGTCTTG	3540
GGGGTTCTCA	ATGGACTGGT	TCTGCTGCCT	GTCTCTTAT	CCTTCTTTGG	ACCGTGTCTT	3600
GAGGTGTCTC	CAGCCAATGG	CCTAAACCGA	CTGCCCACTC	CTTCGCCTGA	GCCGCCTCCA	3660
AGTGTCGTCC	GGTTTGCCGT	GCCTCCTGGT	CACACGAACA	ATGGGTCTGA	TTCTCTCCGAC	3720
TCGGAGTACA	GCTCTCAGAC	CACGGTGTCT	GGCATCAGTG	AGGAGCTCAG	GCAATACGAA	3780
GCACAGCAGG	GTGCCGGAGG	CCCTGCCCCAC	CAAGTGATTG	TGGAAGCCAC	AGAAAACCTT	3840
GTCTTTGCCC	GGTCCACTGT	GGTCCATCCG	GA CTCCAGAC	ATCAGCCTCC	CTTGACCCCTT	3900

CGGCAACAGC CCCACCTGGA CTCTGGCTCC TTGTCCCCTG GACGGCAAGG CCAGCAGCCT 3960  
 CGAAGGGATC CCCCTAGAGA AGGCTTGCGG CCACCCCCCT ACAGACCGCG CAGAGACGCT 4020  
 TTTGAAATTT CTACTGAAGG GCATTCTGGC CCTAGCAATA GGGACCGCTC AGGGCCCCGT 4080  
 GGGGCCCCGT CTCACAACCC TCGGAACCCA ACGTCCACCG CCATGGGCAG CTCTGTGCCC 4140  
 AGCTACTGCC AGCCCATCAC CACTGTGACG GCTTCTGCTT CGGTGACTGT TGCTGTGCAT 4200  
 CCCCCGCTG GACCTGGGCG CAACCCCCGA GGGGGGCCCT GTCCAGGCTA TGAGAGCTAC 4260  
 CCTGAGACTG ATCACGGGGT ATTTGAGGAT CCTCATGTGC CTTTTCATGT CAGGTGTGAG 4320  
 AGGAGGGACT CAAAGGTGGA GGTCATAGAG CTACAGGACG TGGAAATGTGA GGAGAGGCCG 4380  
 TGGGGGAGCA GCTCCAACCTG AGGGTAATTA AAATCTGAAG CAAAGAGGCC AAAGATTGGA 4440  
 AAGCCCCGCC CCCACCTCTT TCCAGAACTG CTTGAAGAGA ACTGCTTGGA ATTATGGGAA 4500  
 GGCAGTTCAT TGTTACTGTA ACTGATTGTA TTATTKKGTG AAATATTTCT ATAAATATTT 4560  
 AARAGGTGTA CACATGTAAT ATACATGGAA ATGCTGTACA GTCTATTTCC TGGGGCCTCT 4620  
 CCACTCCTGC CCCAGAGTGG GGAGACCACA GGGGCCCTTT CCCCTGTGTA CATTGGTCTC 4680  
 TGTGCCACAA CCAAGCTTAA CTTAGTTTTA AAAAAAATCT CCCAGCATAT GTCGCTGCTG 4740  
 CTTAAATATT GTATAATTTA CTTGTATAAT TCTATGCAA TATTGCTTAT GTAATAGGAT 4800  
 TATTTGTAAA GGTTTCTGTT TAAAATATTT TAAATTTGCA TATCACAACC CTGTGGTAGG 4860  
 ATGAATTGTT ACTGTTAACT TTTGAACACG CTATGCGTGG TAATTGTTTA ACGAGCAGAC 4920  
 ATGAAGAAAA CAGGTTAATC CCAGTGGCTT CTCTAGGGGT AGTTGTATAT GGTTGCGATG 4980  
 GGTGGATGTG TGTGTGCATG TGACTTTCCA ATGTACTGTA TTGTGGTTTG TTGTTGTTGT 5040  
 TGCTGTTGTT GTTCATTTTG GTGTTTTTGG TTGCTTTGTA TGATCTTAGC TCTGGCCTAG 5100  
 GTGGGCTGGG AAGGTCCAGG TCTTTTTCTG TCGTGATGCT GGTGGAAAGG TGACCCCAAT 5160  
 CATCTGTCCT ATTCTCTGGG ACTATTC 5187

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1434 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Ala Ser Ala Gly Asn Ala Ala Gly Ala Leu Gly Arg Gln Ala Gly  
 1 5 10 15  
 Gly Gly Arg Arg Arg Arg Thr Gly Gly Pro His Arg Ala Ala Pro Asp

20						25						30					
Arg	Asp	Tyr	Leu	His	Arg	Pro	Ser	Tyr	Cys	Asp	Ala	Ala	Phe	Ala	Leu		
		35					40					45					
Glu	Gln	Ile	Ser	Lys	Gly	Lys	Ala	Thr	Gly	Arg	Lys	Ala	Pro	Leu	Trp		
	50					55					60						
Leu	Arg	Ala	Lys	Phe	Gln	Arg	Leu	Leu	Phe	Lys	Leu	Gly	Cys	Tyr	Ile		
65					70					75					80		
Gln	Lys	Asn	Cys	Gly	Lys	Phe	Leu	Val	Val	Gly	Leu	Leu	Ile	Phe	Gly		
				85					90					95			
Ala	Phe	Ala	Val	Gly	Leu	Lys	Ala	Ala	Asn	Leu	Glu	Thr	Asn	Val	Glu		
			100					105					110				
Glu	Leu	Trp	Val	Glu	Val	Gly	Gly	Arg	Val	Ser	Arg	Glu	Leu	Asn	Tyr		
		115					120					125					
Thr	Arg	Gln	Lys	Ile	Gly	Glu	Glu	Ala	Met	Phe	Asn	Pro	Gln	Leu	Met		
	130					135					140						
Ile	Gln	Thr	Pro	Lys	Glu	Glu	Gly	Ala	Asn	Val	Leu	Thr	Thr	Glu	Ala		
145					150					155					160		
Leu	Leu	Gln	His	Leu	Asp	Ser	Ala	Leu	Gln	Ala	Ser	Arg	Val	His	Val		
				165					170					175			
Tyr	Met	Tyr	Asn	Arg	Gln	Trp	Lys	Leu	Glu	His	Leu	Cys	Tyr	Lys	Ser		
			180					185					190				
Gly	Glu	Leu	Ile	Thr	Glu	Thr	Gly	Tyr	Met	Asp	Gln	Ile	Ile	Glu	Tyr		
		195					200					205					
Leu	Tyr	Pro	Cys	Leu	Ile	Ile	Thr	Pro	Leu	Asp	Cys	Phe	Trp	Glu	Gly		
	210					215					220						
Ala	Lys	Leu	Gln	Ser	Gly	Thr	Ala	Tyr	Leu	Leu	Gly	Lys	Pro	Pro	Leu		
225					230					235					240		
Arg	Trp	Thr	Asn	Phe	Asp	Pro	Leu	Glu	Phe	Leu	Glu	Glu	Leu	Lys	Lys		
				245					250					255			
Ile	Asn	Tyr	Gln	Val	Asp	Ser	Trp	Glu	Glu	Met	Leu	Asn	Lys	Ala	Glu		
			260					265					270				
Val	Gly	His	Gly	Tyr	Met	Asp	Arg	Pro	Cys	Leu	Asn	Pro	Ala	Asp	Pro		
		275					280					285					
Asp	Cys	Pro	Ala	Thr	Ala	Pro	Asn	Lys	Asn	Ser	Thr	Lys	Pro	Leu	Asp		
	290					295					300						
Val	Ala	Leu	Val	Leu	Asn	Gly	Gly	Cys	Gln	Gly	Leu	Ser	Arg	Lys	Tyr		
305					310					315					320		
Met	His	Trp	Gln	Glu	Glu	Leu	Ile	Val	Gly	Gly	Thr	Val	Lys	Asn	Ala		
				325					330					335			
Thr	Gly	Lys	Leu	Val	Ser	Ala	His	Ala	Leu	Gln	Thr	Met	Phe	Gln	Leu		
			340					345					350				
Met	Thr	Pro	Lys	Gln	Met	Tyr	Glu	His	Phe	Arg	Gly	Tyr	Asp	Tyr	Val		
		355					360					365					

Ser His Ile Asn Trp Asn Glu Asp Arg Ala Ala Ala Ile Leu Glu Ala  
 370 375 380  
 Trp Gln Arg Thr Tyr Val Glu Val Val His Gln Ser Val Ala Pro Asn  
 385 390 395 400  
 Ser Thr Gln Lys Val Leu Pro Phe Thr Thr Thr Thr Leu Asp Asp Ile  
 405 410 415  
 Leu Lys Ser Phe Ser Asp Val Ser Val Ile Arg Val Ala Ser Gly Tyr  
 420 425 430  
 Leu Leu Met Leu Ala Tyr Ala Cys Leu Thr Met Leu Arg Trp Asp Cys  
 435 440 445  
 Ser Lys Ser Gln Gly Ala Val Gly Leu Ala Gly Val Leu Leu Val Ala  
 450 455 460  
 Leu Ser Val Ala Ala Gly Leu Gly Leu Cys Ser Leu Ile Gly Ile Ser  
 465 470 475 480  
 Phe Asn Ala Ala Thr Thr Gln Val Leu Pro Phe Leu Ala Leu Gly Val  
 485 490 495  
 Gly Val Asp Asp Val Phe Leu Leu Ala His Ala Phe Ser Glu Thr Gly  
 500 505 510  
 Gln Asn Lys Arg Ile Pro Phe Glu Asp Arg Thr Gly Glu Cys Leu Lys  
 515 520 525  
 Arg Thr Gly Ala Ser Val Ala Leu Thr Ser Ile Ser Asn Val Thr Ala  
 530 535 540  
 Phe Phe Met Ala Ala Leu Ile Pro Ile Pro Ala Leu Arg Ala Phe Ser  
 545 550 555 560  
 Leu Gln Ala Ala Val Val Val Val Phe Asn Phe Ala Met Val Leu Leu  
 565 570 575  
 Ile Phe Pro Ala Ile Leu Ser Met Asp Leu Tyr Arg Arg Glu Asp Arg  
 580 585 590  
 Arg Leu Asp Ile Phe Cys Cys Phe Thr Ser Pro Cys Val Ser Arg Val  
 595 600 605  
 Ile Gln Val Glu Pro Gln Ala Tyr Thr Glu Pro His Ser Asn Thr Arg  
 610 615 620  
 Tyr Ser Pro Pro Pro Tyr Thr Ser His Ser Phe Ala His Glu Thr  
 625 630 635 640  
 His Ile Thr Met Gln Ser Thr Val Gln Leu Arg Thr Glu Tyr Asp Pro  
 645 650 655  
 His Thr His Val Tyr Tyr Thr Thr Ala Glu Pro Arg Ser Glu Ile Ser  
 660 665 670  
 Val Gln Pro Val Thr Val Thr Gln Asp Asn Leu Ser Cys Gln Ser Pro  
 675 680 685  
 Glu Ser Thr Ser Ser Thr Arg Asp Leu Leu Ser Gln Phe Ser Asp Ser  
 690 695 700

001640-0890  
 04T9T580

Ser 705	Leu	His	Cys	Leu	Glu	Pro	Pro	Cys	Thr	Lys	Trp	Thr	Leu	Ser	Ser
					710					715					720
Phe	Ala	Glu	Lys	His	Tyr	Ala	Pro	Phe	Leu	Leu	Lys	Pro	Lys	Ala	Lys
				725					730					735	
Val	Val	Val	Ile	Leu	Leu	Phe	Leu	Gly	Leu	Leu	Gly	Val	Ser	Leu	Tyr
			740					745					750		
Gly	Thr	Thr	Arg	Val	Arg	Asp	Gly	Leu	Asp	Leu	Thr	Asp	Ile	Val	Pro
		755					760					765			
Arg	Glu	Thr	Arg	Glu	Tyr	Asp	Phe	Ile	Ala	Ala	Gln	Phe	Lys	Tyr	Phe
	770					775					780				
Ser	Phe	Tyr	Asn	Met	Tyr	Ile	Val	Thr	Gln	Lys	Ala	Asp	Tyr	Pro	Asn
785					790					795					800
Ile	Gln	His	Leu	Leu	Tyr	Asp	Leu	His	Lys	Ser	Phe	Ser	Asn	Val	Lys
			805						810					815	
Tyr	Val	Met	Leu	Glu	Glu	Asn	Lys	Gln	Leu	Pro	Gln	Met	Trp	Leu	His
			820					825					830		
Tyr	Phe	Arg	Asp	Trp	Leu	Gln	Gly	Leu	Gln	Asp	Ala	Phe	Asp	Ser	Asp
		835					840					845			
Trp	Glu	Thr	Gly	Arg	Ile	Met	Pro	Asn	Asn	Tyr	Lys	Asn	Gly	Ser	Asp
	850					855					860				
Asp	Gly	Val	Leu	Ala	Tyr	Lys	Leu	Leu	Val	Gln	Thr	Gly	Ser	Arg	Asp
865					870					875					880
Lys	Pro	Ile	Asp	Ile	Ser	Gln	Leu	Thr	Lys	Gln	Arg	Leu	Val	Asp	Ala
				885					890					895	
Asp	Gly	Ile	Ile	Asn	Pro	Ser	Ala	Phe	Tyr	Ile	Tyr	Leu	Thr	Ala	Trp
		900						905					910		
Val	Ser	Asn	Asp	Pro	Val	Ala	Tyr	Ala	Ala	Ser	Gln	Ala	Asn	Ile	Arg
		915					920					925			
Pro	His	Arg	Pro	Glu	Trp	Val	His	Asp	Lys	Ala	Asp	Tyr	Met	Pro	Glu
	930					935					940				
Thr	Arg	Leu	Arg	Ile	Pro	Ala	Ala	Glu	Pro	Ile	Glu	Tyr	Ala	Gln	Phe
945					950					955					960
Pro	Phe	Tyr	Leu	Asn	Gly	Leu	Arg	Asp	Thr	Ser	Asp	Phe	Val	Glu	Ala
				965					970					975	
Ile	Glu	Lys	Val	Arg	Val	Ile	Cys	Asn	Asn	Tyr	Thr	Ser	Leu	Gly	Leu
			980					985					990		
Ser	Ser	Tyr	Pro	Asn	Gly	Tyr	Pro	Phe	Leu	Phe	Trp	Glu	Gln	Tyr	Ile
		995					1000					1005			
Ser	Leu	Arg	His	Trp	Leu	Leu	Leu	Ser	Ile	Ser	Val	Val	Leu	Ala	Cys
	1010					1015					1020				
Thr	Phe	Leu	Val	Cys	Ala	Val	Phe	Leu	Leu	Asn	Pro	Trp	Thr	Ala	Gly
1025					1030					1035					1040

Ile Ile Val Met Val Leu Ala Leu Met Thr Val Glu Leu Phe Gly Met  
 1045 1050 1055  
 Met Gly Leu Ile Gly Ile Lys Leu Ser Ala Val Pro Val Val Ile Leu  
 1060 1065 1070  
 Ile Ala Ser Val Gly Ile Gly Val Glu Phe Thr Val His Val Ala Leu  
 1075 1080 1085  
 Ala Phe Leu Thr Ala Ile Gly Asp Lys Asn His Arg Ala Met Leu Ala  
 1090 1095 1100  
 Leu Glu His Met Phe Ala Pro Val Leu Asp Gly Ala Val Ser Thr Leu  
 1105 1110 1115 1120  
 Leu Gly Val Leu Met Leu Ala Gly Ser Glu Phe Asp Phe Ile Val Arg  
 1125 1130 1135  
 Tyr Phe Phe Ala Val Leu Ala Ile Leu Thr Val Leu Gly Val Leu Asn  
 1140 1145 1150  
 Gly Leu Val Leu Leu Pro Val Leu Leu Ser Phe Phe Gly Pro Cys Pro  
 1155 1160 1165  
 Glu Val Ser Pro Ala Asn Gly Leu Asn Arg Leu Pro Thr Pro Ser Pro  
 1170 1175 1180  
 Glu Pro Pro Pro Ser Val Val Arg Phe Ala Val Pro Pro Gly His Thr  
 1185 1190 1195 1200  
 Asn Asn Gly Ser Asp Ser Ser Asp Ser Glu Tyr Ser Ser Gln Thr Thr  
 1205 1210 1215  
 Val Ser Gly Ile Ser Glu Glu Leu Arg Gln Tyr Glu Ala Gln Gln Gly  
 1220 1225 1230  
 Ala Gly Gly Pro Ala His Gln Val Ile Val Glu Ala Thr Glu Asn Pro  
 1235 1240 1245  
 Val Phe Ala Arg Ser Thr Val Val His Pro Asp Ser Arg His Gln Pro  
 1250 1255 1260  
 Pro Leu Thr Pro Arg Gln Gln Pro His Leu Asp Ser Gly Ser Leu Ser  
 1265 1270 1275 1280  
 Pro Gly Arg Gln Gly Gln Gln Pro Arg Arg Asp Pro Pro Arg Glu Gly  
 1285 1290 1295  
 Leu Arg Pro Pro Pro Tyr Arg Pro Arg Arg Asp Ala Phe Glu Ile Ser  
 1300 1305 1310  
 Thr Glu Gly His Ser Gly Pro Ser Asn Arg Asp Arg Ser Gly Pro Arg  
 1315 1320 1325  
 Gly Ala Arg Ser His Asn Pro Arg Asn Pro Thr Ser Thr Ala Met Gly  
 1330 1335 1340  
 Ser Ser Val Pro Ser Tyr Cys Gln Pro Ile Thr Thr Val Thr Ala Ser  
 1345 1350 1355 1360  
 Ala Ser Val Thr Val Ala Val His Pro Pro Pro Gly Pro Gly Arg Asn  
 1365 1370 1375

005440-0034

Pro Arg Gly Gly Pro Cys Pro Gly Tyr Glu Ser Tyr Pro Glu Thr Asp  
 1380 1385 1390

His Gly Val Phe Glu Asp Pro His Val Pro Phe His Val Arg Cys Glu  
 1395 1400 1405

Arg Arg Asp Ser Lys Val Glu Val Ile Glu Leu Gln Asp Val Glu Cys  
 1410 1415 1420

Glu Glu Arg Pro Trp Gly Ser Ser Ser Asn  
 1425 1430

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 11 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Ile Ile Thr Pro Leu Asp Cys Phe Trp Glu Gly  
 1 5 10

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 5 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Leu Ile Val Gly Gly  
 1 5

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 7 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Pro Phe Phe Trp Glu Gln Tyr

"0879166" 14169

1

5

## (2) INFORMATION FOR SEQ ID NO:14:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "primer"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GGACGAATTC AARGTNCAYC ARYTNTGG

28

## (2) INFORMATION FOR SEQ ID NO:15:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "primer"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GGACGAATTC CYTCCCARAA RCANTC

26

## (2) INFORMATION FOR SEQ ID NO:16:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "primer"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GGACGAATTC YTNGANTGYT TYTGGA

27

## (2) INFORMATION FOR SEQ ID NO:17:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

"01979" 01979

(ii) MOLECULE TYPE: other nucleic acid  
 (A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CATACCAGCC AAGCTTGTCTN GGCCARTGCA T

31

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 5288 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GAATTCCGGG GACCGCAAGG AGTGCCGCGG AAGCGCCCGA AGGACAGGCT CGCTCGGCGC	60
GCCGGCTCTC GCTCTTCCGC GAACTGGATG TGGGCAGCGG CGGCCGCAGA GACCTCGGGA	120
CCCCGCGCA ATGTGGCAAT GGAAGGCGCA GGGTCTGACT CCCCggcagc GGCCGCGGCC	180
GCAGCGGCAG CAGCGCCCGC CGTGTGAGCA GCAGCAGCGG CTGGTCTGTC AACCGGAGCC	240
CGAGCCCGAG CAGCCTGCGG CCAGCAGCGT CCTCGCAAGC CGAGCGCCCA GGCGCGCCAG	300
GAGCCCGCAG CAGCGGCAGC AGCGCGCCGG GCCGCCCGGG AAGCCTCCGT CCCCgCGGCG	360
GCGGCGGCGG CGGCGGCGGC AACATGGCCT CGGCTGGTAA CGCCGCCGAG CCCCAGGACC	420
GCGGCGGCGG CGGCAGCGGC TGTATCGGTG CCCCgggacg GCCGGCTGGA GGCGGGAGGC	480
GCAGACGGAC GGGGGGGCTG CGCCGTGCTG CCGCGCCGGA CCGGGACTAT CTGCACCGGC	540
CCAGCTACTG CGACGCCGCC TTCGCTCTGG AGCAGATTTC CAAGGGGAAG GCTACTGGCC	600
GGAAAGCGCC ACTGTGGCTG AGAGCGAAGT TTCAGAGACT CTTATTTAAA CTGGGTGTGT	660
ACATTCAAAA AAAGTGCAGC AAGTTCTTGG TTGTGGGCCT CCTCATATTT GGGGCCTTCG	720
CGGTGGGATT AAAAGCAGCG AACCTCGAGA CCAACGTGGA GGAGCTGTGG GTGGAAGTTG	780
GAGGACGAGT AAGTCGTGAA TTAAATTATA CTCGCCAGAA GATTGGAGAA GAGGCTATGT	840
TTAATCCTCA ACTCATGATA CAGACCCCTA AAGAAGAAGG TGCTAATGTC CTGACCACAG	900
AAGCGCTCCT ACAACACCTG GACTCGGCAC TCCAGGCCAG CCGTGTCCAT GTATACATGT	960
ACAACAGGCA GTGGAAATTG GAACATTTGT GTTACAAATC AGGAGAGCTT ATCACAGAAA	1020
CAGGTTACAT GGATCAGATA ATAGAATATC TTTACCCTTG TTTGATTATT ACACCTTTGG	1080
ACTGCTTCTG GGAAGGGGCG AAATTACAGT CTGGGACAGC ATACCTCCTA GGTAACCTC	1140

0016140-0016140

CTTTGCGGTG	GACAAACTTC	GACCCTTTGG	AATTCCTGGA	AGAGTTAAAG	AAAATAAACT	1200
ATCAAGTGGA	CAGCTGGGAG	GAAATGCTGA	ATAAGGCTGA	GGTTGGTCAT	GGTTACATGG	1260
ACCGCCCCTG	CCTCAATCCG	GCCGATCCAG	ACTGCCCCGC	CACAGCCCCC	AACAAAAATT	1320
CAACCAAACC	TCTTGATATG	GCCCTTGTTT	TGAATGGTGG	ATGTCATGGC	TTATCCAGAA	1380
AGTATATGCA	CTGGCAGGAG	GAGTTGATTG	TGGGTGGCAC	AGTCAAGAAC	AGCACTGGAA	1440
AACTCGTCAG	CGCCCATGCC	CTGCAGACCA	TGTTCCAGTT	AATGACTCCC	AAGCAAATGT	1500
ACGAGCACTT	CAAGGGGTAC	GAGTATGTCT	CACACATCAA	CTGGAACGAG	GACAAAGCGG	1560
CAGCCATCCT	GGAGGCCTGG	CAGAGGACAT	ATGTGGAGGT	GGTTCATCAG	AGTGTGCAC	1620
AGAACTCCAC	TCAAAGGTG	CTTTCCTTCA	CCACCACGAC	CCTGGACGAC	ATCCTGAAAT	1680
CCTTCTCTGA	CGTCAGTGTC	ATCCGCGTGG	CCAGCGGCTA	CTTACTCATG	CTCGCCTATG	1740
CCTGTCTAAC	CATGCTGCGC	TGGGACTGCT	CCAAGTCCCA	GGGTGCCGTG	GGGCTGGCTG	1800
GGCTCCTGCT	GGTTGCACTG	TCAGTGGCTG	CAGGACTGGG	CCTGTGCTCA	TTGATCGGAA	1860
TTTCCTTTAA	CGCTGCAACA	ACTCAGGTTT	TGCCATTTCT	CGCTCTTGGT	GTTGGTGTGG	1920
ATGATGTTTT	TCTTCTGGCC	CACGCCTTCA	GTGAAACAGG	ACAGAATAAA	AGAATCCCTT	1980
TTGAGGACAG	GACCGGGGAG	TGCCTGAAGC	GCACAGGAGC	CAGCGTGGCC	CTCACGTCCA	2040
TCAGCAATGT	CACAGCCTTC	TTCATGGCCG	CGTTAATCCC	AATTCCCCTG	CTGCGGGCGT	2100
TCTCCCTCCA	GGCAGCGGTA	GTAGTGGTGT	TCAATTTTGC	CATGGTTCTG	CTCATTTTTT	2160
CTGCAATTCT	CAGCATGGAT	TTATATCGAC	GCGAGGACAG	GAGACTGGAT	ATTTTCTGCT	2220
GTTTTACAAG	CCCCTGCGTC	AGCAGAGTGA	TTCAGGTTGA	ACCTCAGGCC	TACACCGACA	2280
CACACGACAA	TACCCGCTAC	AGCCCCCAC	CTCCCTACAG	CAGCCACAGC	TTTGCCCATG	2340
AAACGCAGAT	TACCATGCAG	TCCACTGTCC	AGCTCCGCAC	GGAGTACGAC	CCCCACACGC	2400
ACGTGTACTA	CACCACCGCT	GAGCCGCGCT	CCGAGATCTC	TGTGCAGCCC	GTCACCGTGA	2460
CACAGGACAC	CCTCAGCTGC	CAGAGCCCAG	AGAGCACCAG	CTCCACAAGG	GACCTGCTCT	2520
CCCAGTTCTC	CGACTCCAGC	CTCCACTGCC	TCGAGCCCCC	CTGTACGAAG	TGGACACTCT	2580
CATCTTTTGC	TGAGAAGCAC	TATGCTCCTT	TCCTCTTGAA	ACCAAAGCC	AAGGTAGTGG	2640
TGATCTTCCT	TTTTCTGGGC	TTGCTGGGGG	TCAGCCTTTA	TGGCACCACC	CGAGTGAGAG	2700
ACGGGCTGGA	CCTTACGGAC	ATTGTACCTC	GGGAAACCAG	AGAATATGAC	TTTATTGCTG	2760
CACAATTCAA	ATACTTTTCT	TTCTACAACA	TGTATATAGT	CACCCAGAAA	GCAGACTACC	2820
CGAATATCCA	GCACTTACTT	TACGACCTAC	ACAGGAGTTT	CAGTAACGTG	AAGTATGTCA	2880
TGTTGGAAGA	AAACAAACAG	CTTCCCAAAA	TGTGGCTGCA	CTACTTCAGA	GACTGGCTTC	2940
AGTCACTTCA	GGATGCATTT	GACAGTGACT	GGGAAACCGG	GAAAATCATG	CCAAACAATT	3000
ACAAGAATGG	ATCAGACGAT	GGAGTCCTTG	CCTACAAACT	CCTGGTGCAA	ACCGGCAGCC	3060

GCGATAAGCC	CATCGACATC	AGCCAGTTGA	CTAAACAGCG	TCTGGTGGAT	GCAGATGGCA	3120
TCATTAATCC	CAGCGCTTTC	TACATCTACC	TGACGGCTTG	GGTCAGCAAC	GACCCCGTCG	3180
CGTATGCTGC	CTCCCAGGCC	AACATCCGGC	CACACCGACC	AGAATGGGTC	CACGACAAAG	3240
CCGACTACAT	GCCTGAAACA	AGGCTGAGAA	TCCCGGCAGC	AGAGCCCATC	GAGTATGCCC	3300
AGTTCCCTTT	CTACCTCAAC	GGGTTGCGGG	ACACCTCAGA	CTTTGTGGAG	GCAATTGAAA	3360
AAGTAAGGAC	CATCTGCAGC	AACTATACGA	GCCTGGGGCT	GTCCAGTTAC	CCCAACGGCT	3420
ACCCCTTCCT	CTTCTGGGAG	CAGTACATCG	GCCTCCGCCA	CTGGCTGCTG	CTGTTCATCA	3480
GCGTGGTGTT	GGCCTGCACA	TTCCTCGTGT	GCGCTGTCTT	CCTTCTGAAC	CCCTGGACGG	3540
CCGGGATCAT	TGTGATGGTC	CTGGCGCTGA	TGACGGTCGA	GCTGTTCCGC	ATGATGGGCC	3600
TCATCGGAAT	CAAGCTCAGT	GCCGTGCCCC	TGGTCATCCT	GATCGCTTCT	GTTGGCATAG	3660
GAGTGGAGTT	CACCGTTCAC	GTTGCTTTGG	CCTTTCTGAC	GGCCATCGGC	GACAAGAACC	3720
GCAGGGCTGT	GCTTGCCCTG	GAGCACATGT	TTGCACCCGT	CCTGGATGGC	GCCGTGTCCA	3780
CTCTGCTGGG	AGTGCTGATG	CTGGCGGGAT	CTGAGTTCGA	CTTCATTGTC	AGGTATTTCT	3840
TTGCTGTGCT	GGCGATCCTC	ACCATCCTCG	GCGTTCTCAA	TGGGCTGGTT	TTGCTTCCCG	3900
TGCTTTTGTG	TTTCTTTGGA	CCATATCCTG	AGGTGTCTCC	AGCCAACGGC	TTGAACCGCC	3960
TGCCCACACC	CTCCCCTGAG	CCACCCCCCA	GCGTGGTCCG	CTTCGCCATG	CCGCCCCGCC	4020
ACACGCACAG	CGGGTCTGAT	TCCTCCGACT	CGGAGTATAG	TTCCCAGACG	ACAGTGTGAG	4080
GCCTCAGCGA	GGAGCTTCGG	CACTACGAGG	CCCAGCAGGG	CGCGGGAGGC	CCTGCCCCACC	4140
AAGTGATCGT	GGAAGCCACA	GAAAACCCCG	TCTTCGCCCA	CTCCACTGTG	GTCCATCCCG	4200
AATCCAGGCA	TCACCCACCC	TCAACCCCGA	GACAGCAGCC	CCACCTGGAC	TCAGGGTCCC	4260
TGCCTCCCGG	ACGGCAAGGC	CAGCAGCCCC	GCAGGGACCC	CCCCAGAGAA	GGCTTGTGGC	4320
CACCCCTCTA	CAGACCGCGC	AGAGACGCTT	TTGAAATTTT	TACTGAAGGG	CATTCTGGCC	4380
CTAGCAATAG	GGCCCGCTGG	GGCCCTCGCG	GGGCCCCGTT	TCACAACCCT	CGGAACCCAG	4440
CGTCCACTGC	CATGGGCAGC	TCCGTGCCCC	GCTACTGCCA	GCCCATCACC	ACTGTGACGG	4500
CTTCTGCCTC	CGTGAAGTGC	GCCGTGCACC	CGCCGCCTGT	CCCTGGGCCT	GGGCGGAACC	4560
CCCGAGGGGG	ACTCTGCCCA	GGCTACCCTG	AGACTGACCA	CGGCCTGTTT	GAGGACCCCC	4620
ACGTGCCTTT	CCACGTCCGG	TGTGAGAGGA	GGGATTGCAA	GGTGAAGTCT	ATTGAGCTGC	4680
AGGACGTGGA	ATGCGAGGAG	AGGCCCCGGG	GAAGCAGCTC	CAACTGAGGG	TGATTAAAAT	4740
CTGAAGCAAA	GAGGCCAAAG	ATTGGAAACC	CCCCACCCCC	ACCTCTTTCC	AGAACTGCTT	4800
GAAGAGAACT	GGTTGGAGTT	ATGGAAAAGA	TGCCCTGTGC	CAGGACAGCA	GTTTCATTGT	4860
ACTGTAACCG	ATTGTATTAT	TTTGTTAAAT	ATTTCTATAA	ATATTTAAGA	GATGTACACA	4920

(2) INFORMATION FOR SEO ID NO:19:

(A) LENGTH: 1447 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Met 1	Ala	Ser	Ala	Gly 5	Asn	Ala	Ala	Glu	Pro 10	Gln	Asp	Arg	Gly	Gly 15	Gly
Gly	Ser	Gly	Cys 20	Ile	Gly	Ala	Pro	Gly 25	Arg	Pro	Ala	Gly	Gly 30	Gly	Arg
Arg	Arg	Arg 35	Thr	Gly	Gly	Leu	Arg 40	Arg	Ala	Ala	Ala	Pro 45	Asp	Arg	Asp
Tyr	Leu 50	His	Arg	Pro	Ser	Tyr 55	Cys	Asp	Ala	Ala	Phe 60	Ala	Leu	Glu	Gln
Ile 65	Ser	Lys	Gly	Lys 70	Ala	Thr	Gly	Arg	Lys	Ala 75	Pro	Leu	Trp	Leu 80	Arg
Ala	Lys	Phe	Gln	Arg 85	Leu	Leu	Phe	Lys	Leu 90	Gly	Cys	Tyr	Ile	Gln 95	Lys
Asn	Cys	Gly	Lys 100	Phe	Leu	Val	Val	Gly 105	Leu	Leu	Ile	Phe	Gly 110	Ala	Phe
Ala	Val	Gly 115	Leu	Lys	Ala	Ala	Asn 120	Leu	Glu	Thr	Asn	Val 125	Glu	Glu	Leu
Trp 130	Val	Glu	Val	Gly	Gly	Arg 135	Val	Ser	Arg	Glu	Leu 140	Asn	Tyr	Thr	Arg
Gln 145	Lys	Ile	Gly	Glu 150	Glu	Ala	Met	Phe	Asn 155	Pro	Gln	Leu	Met	Ile	Gln 160
Thr	Pro	Lys	Glu	Glu 165	Gly	Ala	Asn	Val	Leu 170	Thr	Thr	Glu	Ala	Leu 175	Leu
Gln	His	Leu	Asp	Ser	Ala	Leu	Gln	Ala	Ser	Arg	Val	His	Val	Tyr	Met

180					185					190					
Tyr	Asn	Arg	Gln	Trp	Lys	Leu	Glu	His	Leu	Cys	Tyr	Lys	Ser	Gly	Glu
	195						200					205			
Leu	Ile	Thr	Glu	Thr	Gly	Tyr	Met	Asp	Gln	Ile	Ile	Glu	Tyr	Leu	Tyr
	210					215					220				
Pro	Cys	Leu	Ile	Ile	Thr	Pro	Leu	Asp	Cys	Phe	Trp	Glu	Gly	Ala	Lys
225					230					235					240
Leu	Gln	Ser	Gly	Thr	Ala	Tyr	Leu	Leu	Gly	Lys	Pro	Pro	Leu	Arg	Trp
				245					250					255	
Thr	Asn	Phe	Asp	Pro	Leu	Glu	Phe	Leu	Glu	Glu	Leu	Lys	Lys	Ile	Asn
			260					265					270		
Tyr	Gln	Val	Asp	Ser	Trp	Glu	Glu	Met	Leu	Asn	Lys	Ala	Glu	Val	Gly
		275					280					285			
His	Gly	Tyr	Met	Asp	Arg	Pro	Cys	Leu	Asn	Pro	Ala	Asp	Pro	Asp	Cys
	290					295					300				
Pro	Ala	Thr	Ala	Pro	Asn	Lys	Asn	Ser	Thr	Lys	Pro	Leu	Asp	Met	Ala
305						310					315				320
Leu	Val	Leu	Asn	Gly	Gly	Cys	His	Gly	Leu	Ser	Arg	Lys	Tyr	Met	His
				325					330					335	
Trp	Gln	Glu	Glu	Leu	Ile	Val	Gly	Gly	Thr	Val	Lys	Asn	Ser	Thr	Gly
			340					345					350		
Lys	Leu	Val	Ser	Ala	His	Ala	Leu	Gln	Thr	Met	Phe	Gln	Leu	Met	Thr
		355					360					365			
Pro	Lys	Gln	Met	Tyr	Glu	His	Phe	Lys	Gly	Tyr	Glu	Tyr	Val	Ser	His
	370					375					380				
Ile	Asn	Trp	Asn	Glu	Asp	Lys	Ala	Ala	Ala	Ile	Leu	Glu	Ala	Trp	Gln
385						390					395				400
Arg	Thr	Tyr	Val	Glu	Val	Val	His	Gln	Ser	Val	Ala	Gln	Asn	Ser	Thr
				405					410					415	
Gln	Lys	Val	Leu	Ser	Phe	Thr	Thr	Thr	Thr	Leu	Asp	Asp	Ile	Leu	Lys
			420					425					430		
Ser	Phe	Ser	Asp	Val	Ser	Val	Ile	Arg	Val	Ala	Ser	Gly	Tyr	Leu	Leu
		435					440					445			
Met	Leu	Ala	Tyr	Ala	Cys	Leu	Thr	Met	Leu	Arg	Trp	Asp	Cys	Ser	Lys
	450					455					460				
Ser	Gln	Gly	Ala	Val	Gly	Leu	Ala	Gly	Val	Leu	Leu	Val	Ala	Leu	Ser
465						470					475			480	
Val	Ala	Ala	Gly	Leu	Gly	Leu	Cys	Ser	Leu	Ile	Gly	Ile	Ser	Phe	Asn
				485					490					495	
Ala	Ala	Thr	Thr	Gln	Val	Leu	Pro	Phe	Leu	Ala	Leu	Gly	Val	Gly	Val
			500					505					510		
Asp	Asp	Val	Phe	Leu	Leu	Ala	His	Ala	Phe	Ser	Glu	Thr	Gly	Gln	Asn

08916140-08916140

515					520					525						
Lys	Arg	Ile	Pro	Phe	Glu	Asp	Arg	Thr	Gly	Glu	Cys	Leu	Lys	Arg	Thr	
530					535					540						
Gly	Ala	Ser	Val	Ala	Leu	Thr	Ser	Ile	Ser	Asn	Val	Thr	Ala	Phe	Phe	
545					550					555					560	
Met	Ala	Ala	Leu	Ile	Pro	Ile	Pro	Ala	Leu	Arg	Ala	Phe	Ser	Leu	Gln	
565					570					575						
Ala	Ala	Val	Val	Val	Val	Phe	Asn	Phe	Ala	Met	Val	Leu	Leu	Ile	Phe	
580					585					590						
Pro	Ala	Ile	Leu	Ser	Met	Asp	Leu	Tyr	Arg	Arg	Glu	Asp	Arg	Arg	Leu	
595					600					605						
Asp	Ile	Phe	Cys	Cys	Phe	Thr	Ser	Pro	Cys	Val	Ser	Arg	Val	Ile	Gln	
610					615					620						
Val	Glu	Pro	Gln	Ala	Tyr	Thr	Asp	Thr	His	Asp	Asn	Thr	Arg	Tyr	Ser	
625					630					635					640	
Pro	Pro	Pro	Pro	Tyr	Ser	Ser	His	Ser	Phe	Ala	His	Glu	Thr	Gln	Ile	
645					650					655						
Thr	Met	Gln	Ser	Thr	Val	Gln	Leu	Arg	Thr	Glu	Tyr	Asp	Pro	His	Thr	
660					665					670						
His	Val	Tyr	Tyr	Thr	Thr	Ala	Glu	Pro	Arg	Ser	Glu	Ile	Ser	Val	Gln	
675					680					685						
Pro	Val	Thr	Val	Thr	Gln	Asp	Thr	Leu	Ser	Cys	Gln	Ser	Pro	Glu	Ser	
690					695					700						
Thr	Ser	Ser	Thr	Arg	Asp	Leu	Leu	Ser	Gln	Phe	Ser	Asp	Ser	Ser	Leu	
705					710					715					720	
His	Cys	Leu	Glu	Pro	Pro	Cys	Thr	Lys	Trp	Thr	Leu	Ser	Ser	Phe	Ala	
725					730					735						
Glu	Lys	His	Tyr	Ala	Pro	Phe	Leu	Leu	Lys	Pro	Lys	Ala	Lys	Val	Val	
740					745					750						
Val	Ile	Phe	Leu	Phe	Leu	Gly	Leu	Leu	Gly	Val	Ser	Leu	Tyr	Gly	Thr	
755					760					765						
Thr	Arg	Val	Arg	Asp	Gly	Leu	Asp	Leu	Thr	Asp	Ile	Val	Pro	Arg	Glu	
770					775					780						
Thr	Arg	Glu	Tyr	Asp	Phe	Ile	Ala	Ala	Gln	Phe	Lys	Tyr	Phe	Ser	Phe	
785					790					795					800	
Tyr	Asn	Met	Tyr	Ile	Val	Thr	Gln	Lys	Ala	Asp	Tyr	Pro	Asn	Ile	Gln	
805					810					815						
His	Leu	Leu	Tyr	Asp	Leu	His	Arg	Ser	Phe	Ser	Asn	Val	Lys	Tyr	Val	
820					825					830						
Met	Leu	Glu	Glu	Asn	Lys	Gln	Leu	Pro	Lys	Met	Trp	Leu	His	Tyr	Phe	
835					840					845						
Arg	Asp	Trp	Leu	Gln	Gly	Leu	Gln	Asp	Ala	Phe	Asp	Ser	Asp	Trp	Glu	
850					855					860						

00015140-000107

Thr 865	Gly	Lys	Ile	Met	Pro 870	Asn	Asn	Tyr	Lys	Asn 875	Gly	Ser	Asp	Asp	Gly 880
Val	Leu	Ala	Tyr	Lys 885	Leu	Leu	Val	Gln	Thr 890	Gly	Ser	Arg	Asp	Lys 895	Pro
Ile	Asp	Ile	Ser 900	Gln	Leu	Thr	Lys	Gln 905	Arg	Leu	Val	Asp	Ala 910	Asp	Gly
Ile	Ile	Asn 915	Pro	Ser	Ala	Phe	Tyr 920	Ile	Tyr	Leu	Thr	Ala 925	Trp	Val	Ser
Asn	Asp 930	Pro	Val	Ala	Tyr	Ala 935	Ala	Ser	Gln	Ala	Asn 940	Ile	Arg	Pro	His
Arg 945	Pro	Glu	Trp	Val	His 950	Asp	Lys	Ala	Asp	Tyr 955	Met	Pro	Glu	Thr	Arg 960
Leu	Arg	Ile	Pro	Ala 965	Ala	Glu	Pro	Ile	Glu 970	Tyr	Ala	Gln	Phe	Pro 975	Phe
Tyr	Leu	Asn 980	Gly	Leu	Arg	Asp	Thr	Ser 985	Asp	Phe	Val	Glu	Ala 990	Ile	Glu
Lys	Val 995	Arg	Thr	Ile	Cys	Ser	Asn 1000	Tyr	Thr	Ser	Leu	Gly 1005	Leu	Ser	Ser
Tyr 1010	Pro	Asn	Gly	Tyr	Pro	Phe 1015	Leu	Phe	Trp	Glu	Gln 1020	Tyr	Ile	Gly	Leu
Arg 1025	His	Trp	Leu	Leu	Leu 1030	Phe	Ile	Ser	Val	Val 1035	Leu	Ala	Cys	Thr	Phe 1040
Leu	Val	Cys	Ala	Val 1045	Phe	Leu	Leu	Asn	Pro 1050	Trp	Thr	Ala	Gly	Ile 1055	Ile
Val	Met	Val 1060	Leu	Ala	Leu	Met	Thr	Val 1065	Glu	Leu	Phe	Gly	Met 1070	Met	Gly
Leu	Ile	Gly 1075	Ile	Lys	Leu	Ser	Ala 1080	Val	Pro	Val	Val	Ile 1085	Leu	Ile	Ala
Ser 1090	Val	Gly	Ile	Gly	Val	Glu 1095	Phe	Thr	Val	His 1100	Val	Ala	Leu	Ala	Phe
Leu 1105	Thr	Ala	Ile	Gly	Asp 1110	Lys	Asn	Arg	Arg	Ala 1115	Val	Leu	Ala	Leu	Glu 1120
His	Met	Phe	Ala 1125	Pro	Val	Leu	Asp	Gly	Ala 1130	Val	Ser	Thr	Leu 1135	Leu	Gly
Val	Leu	Met 1140	Leu	Ala	Gly	Ser	Glu 1145	Phe	Asp	Phe	Ile	Val 1150	Arg	Tyr	Phe
Phe	Ala 1155	Val	Leu	Ala	Ile	Leu	Thr 1160	Ile	Leu	Gly	Val	Leu 1165	Asn	Gly	Leu
Val 1170	Leu	Leu	Pro	Val	Leu	Leu 1175	Ser	Phe	Phe	Gly	Pro 1180	Tyr	Pro	Glu	Val
Ser 1185	Pro	Ala	Asn	Gly	Leu 1190	Asn	Arg	Leu	Pro	Thr 1195	Pro	Ser	Pro	Glu	Pro 1200

[illegible]

## WHAT IS CLAIMED IS:

1. An assay for phenotyping the *patched* status of a cell, comprising detecting, in a sample of mammalian cells, the presence or absence of a genetic lesion characterized by at least one of (i) aberrant modification or mutation of a *patched* gene, and (ii) mis-expression of said *patched* gene.
2. The assay of claim 1, wherein detecting said lesion includes:
  - i. providing a diagnostic probe comprising a nucleic acid including a region of nucleotide sequence which hybridizes to a sense or antisense sequence of said *patched* gene, or naturally occurring mutants thereof, or 5' or 3' flanking sequences naturally associated with said gene;
  - ii. combining said probe with nucleic acid from said cell sample; and
  - iii. detecting, by hybridization of said probe to said cellular nucleic acid, the existence of at least one of a deletion of one or more nucleotides from said *patched* gene, an addition of one or more nucleotides to said *patched* gene, a substitution of one or more nucleotides of said *patched* gene, a gross chromosomal rearrangement of all or a portion of said *patched* gene, a gross alteration in the level of an mRNA transcript of said *patched* gene, or a non-wild type splicing pattern of an mRNA transcript of said *patched* gene.
3. The assay of claim 2, wherein hybridization of said probe further comprises subjecting the probe and cellular nucleic acid to a polymerase chain reaction (PCR) and detecting abnormalities in an amplified product.
4. The assay of claim 2, wherein said probe hybridizes under stringent conditions to a nucleic acid designated by SEQ ID No. 9 or 18.
5. The assay of claim 2, wherein said probe hybridizes under stringent conditions to a nucleic acid designated by SEQ ID No. 18.
6. The assay of claim 2, wherein said probe further comprises a label group attached to said nucleic acid and able to be detected.
7. The assay of claim 1, wherein detecting said lesion comprises ascertaining, from a methylation pattern of said *patched* gene, the presence or absence of aberrant methylation of said *patched* gene.
8. The assay of claim 7, wherein the methylation pattern of said *patched* gene is determined by combining nucleic acid of said cell sample with one or more methylation-sensitive restriction endonucleases and determining the restriction digest pattern of at least a portion of said *patched* gene.

9. The assay of claim 1, wherein detecting said lesion comprises detecting the presence or absence of a non-wild type level of a *patched* protein product of said *patched* gene in cells of said cell sample.
10. The assay of claim 9, wherein the level of said *patched* protein is detected in an immunoassay.
11. The assay claim 1, wherein detecting said lesion comprises ascertaining, relative to a wild-type level of *hedgehog*-dependent *patched* signal transduction, the ability of cells in said cell sample to respond to *hedgehog* induction.
12. The assay of claim 1, wherein said cell sample is obtained from a human patient.
13. A method for diagnosing a genetic predisposition of an animal for at least one of a developmental abnormality or a proliferative disorder marked by abberant expression or activity of a *patched* gene or gene product, the method comprising detecting the presence of a predisposing mutation in a *patched* gene in cells of said animal, wherein the presence of said predisposing mutation indicates that said individual has a genetic predisposition for at least one of developmental abnormalities or a proliferative disorder.
14. The method of claim 13, wherein said genetic predisposition is basal cell nevus syndrome.
15. The method of claim 13, wherein said genetic predisposition is a predisposition for developing a carcinoma
16. The method of claim 13, wherein said genetic predisposition is a predisposition for developing a meningiomas.
17. The method of claim 13, wherein said genetic predisposition is a predisposition for developing a medullomas
18. The method of claim 13, wherein said genetic predisposition is a predisposition for developing a fibroma.
19. The method of claim 13, wherein said detecting step comprises analyzing a nucleic acid sample obtained from said animal.
20. The method of claim 13, wherein said detecting step comprises functional analysis of *patched* protein function.
21. The method of claim 13, wherein said detecting step comprises detecting antibody binding to abnormal *patched* protein.
22. A method for characterizing the phenotype of a tumor, comprising detecting the presence of an oncogenic *patched* mutation in cells of the tumor, wherein the presence

of said oncogenic mutation indicates that said tumor has a patched-associated phenotype.

20. The method of claim 19, wherein said tumor is a carcinoma.
21. The method of claim 20, wherein said carcinoma is a basal cell carcinoma.
- 5 22. The method of claim 19, wherein said tumor is a meningioma.
23. The method of claim 19, wherein said tumor is a medulloma
24. The method of claim 19, wherein said tumor is a fibroma.
25. The method of claim 19, wherein said oncogenic *patched* mutation are detected by analyzing DNA of said tumor.
- 10 26. The method of claim 19, wherein said oncogenic *patched* mutation are detected by mRNA of said tumor.
27. The method of claim 19, wherein said detecting step comprises functional analysis of patched protein function.
- 15 28. The method of claim 19, wherein said detecting step comprises detecting antibody binding to abnormal patched protein.
29. A genetically engineered mammalian cell predisposed to develop a proliferative phenotype as a result of transfection of said mammalian cell with at least one nucleic acid construct which inhibits expression of an endogenous *patched* gene or alters the signal transduction activity of a wild-type *patched* protein.
- 20 30. The cell of claim 26, wherein the cell develops a carcinoma phenotype.
31. The cell of claim 30, wherein the cell develops a basal cell carcinoma phenotype.
32. The cell of claim 26, wherein the cell develops a meningioma phenotype.
33. The cell of claim 26, wherein the cell develops a medulloma phenotype.
34. The cell of claim 26, wherein the cell develops a fibroma phenotype.
- 25 35. A method for treating an animal having a disorder characterized by loss-of-function of a *patched* gene, comprising transfecting cells of the animal with an expression construct encoding a *patched* protein.
36. The method of claim 35, wherein the cells are transfected *in vivo*.
37. The method of claim 35, wherein the cells are transfected *in vitro*.
- 30 38. The method of claim 35, wherein the expression construct is a viral vector.
39. The method of claim 35, wherein the transfected cells include epithelial cells.

40. The method of claim 35, wherein the transfected cells include neuronal cells.
41. The method of claim 35, wherein the transfected cells include carcinoma cells.
42. The method of claim 41, wherein the carcinoma cells are basal cell carcinoma cells.
43. The method of claim 35, wherein the transfected cells include meningioma cells.
- 5 43. The method of claim 35, wherein the transfected cells include medulloma cells.
44. The method of claim 35, wherein the transfected cells include fibroma cells.
45. A method for treating an animal having a disorder characterized by loss-of-function of a *patched* gene, comprising administering to the animal an agent which inhibits derepression of one or more *patched*-dependent genes.

*PATCHED GENES AND THEIR USES*

## ABSTRACT OF THE INVENTION

Methods for isolating patched genes, particularly mammalian patched genes, including the mouse and human *patched* genes, as well as invertebrate patched genes and sequences, are provided. Decreased expression of *patched* is associated with the occurrence of human cancers, particularly basal cell carcinomas of the skin. The cancers may be familial, having as a component of risk an inherited genetic predisposition, or may be sporadic. The *patched* and *hedgehog* genes are useful in creating transgenic animal models for these human cancers. The *patched* nucleic acid compositions find use in identifying homologous or related proteins and the DNA sequences encoding such proteins; in producing compositions that modulate the expression or function of the protein; and in studying associated physiological pathways. In addition, modulation of the gene activity *in vivo* is used for prophylactic and therapeutic purposes, such as treatment of cancer, identification of cell type based on expression, and the like. The DNA is further used as a diagnostic for a genetic predisposition to cancer, and to identify specific cancers having mutations in this gene.





FIGURE 1

FIGURE 2



08946440-08949

Figure 3




Figure 5




Figure 4




Figure 6






Figure 7



Declaration, Petition and Power of Attorney For Patent Application

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled:

***PATCHED GENES AND USES RELATED THERETO***

the specification of which is filed herewith in the U.S. Patent and Trademark Office.

I do not know and do not believe that the subject matter of this application was ever known or used in the United States before my invention thereof or patented or described in any printed publication in any country before my invention thereof or more than one year prior to the date of this application, and that said subject matter has not been patented or made the subject of an issued inventor's certificate in any country foreign to the United States on an application filed by me or my legal representatives or assigns more than twelve months prior to the date of this application; that I acknowledge my duty to disclose information of which I am aware which is material to the examination of this application, that no application for patent or inventor's certificate on the subject matter of this application has been filed by me or my representatives or assigns in any country foreign to the United States, except those identified below, and that I have reviewed and understand the contents of the specification, including the claims as amended by any amendment referred to herein.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

00916140-000197

## PRIORITY CLAIM

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed.

Check one: ☒ no such applications have been filed.  
☐ such applications have been filed as follows

EARLIEST FOREIGN APPLICATION(S), IF ANY FILED WITHIN 12 MONTHS  
(6 MONTHS FOR DESIGN) PRIOR TO THIS U.S. APPLICATION

Country	Application Number	Date of Filing (month,day,year)	Priority Claimed Under 35 USC 119
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No
			<input type="checkbox"/> Yes <input type="checkbox"/> No

(6 MONTHS FOR DESIGN) PRIOR TO THIS U.S. APPLICATION


45 T 2330 04T9T630

CLAIM FOR BENEFIT OF EARLIER U.S./PCT APPLICATION(S)

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application. As to subject matter of this application which is common to my earlier United States application, if any, described below, I do not believe that the same was ever known or used in the United States before my invention thereof or patented or described in any printed publication in any country before my invention thereof or more than one year prior to said earlier application, or in public use or on sale in the United States more than one year prior to said earlier application, that the said common subject matter has not been patented or made the subject of an inventor's certificate issued before the date of said earlier application in any country foreign to the United States on an application, filed by me or my legal representatives or assigns more than twelve months prior to said application and that no application for patent or inventor's certificate on said subject matter has been filed by me or my representatives or assigns in any country foreign to the United States except those identified herein.

08/656,055  
(Application Serial No.)

31 May 1996  
(Filing Date)

PENDING  
(Status)  
(patented, pending, aband.)

08/540,406  
(Application Serial No.)

06 October 1995  
(Filing Date)

PENDING  
(Status)  
(patented, pending, aband.)

08/319,745  
(Application Serial No.)

07 October 1994  
(Filing Date)

ABANDONED  
(Status)  
(patented, pending, aband.)

267280-047680

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorneys and/or agents to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

Donald W. Muirhead	Reg. No. 33,978	Charles H. Cella	Reg. No. 38,099
Beth E. Arnold	Reg. No. 35,430	John C. Gorecki	Reg. No. 38,741
Matthew P. Vincent	Reg. No. 36,709	Edward J. Kelly	Reg. No. 38,936
		Sayoko Blodgett-Ford	Reg. No. 40,516

Send Correspondence to: Patent Group  
Foley, Hoag & Eliot LLP  
One Post Office Square  
Boston, MA 02109

Direct Telephone Calls to: Matthew P. Vincent, Ph.D. (617) 832-1000

Wherefore I petition that letters patent be granted to me for the invention or discovery described and claimed in the attached specification and claims, and hereby subscribe my name to said specification and claims and to the foregoing declaration, power of attorney, and this petition.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor Matthew P. Scott	
Inventor's signature	Date
Residence 914 Wing Place, Stanford, CA 94035	
Citizenship USA	
Post Office Address (if different)	

Full name of second inventor Lisa V. Goodrich	
Inventor's signature	Date
Residence 66 Newell Road, Palo Alto, CA 94303	
Citizenship USA	
Post Office Address (if different)	

Full name of third inventor

Ronald L. Johnson

Inventor's signature

Date

Residence

1528 Hudson Street, Apt. 7, Redwood City, CA 94061

Citizenship

USA

Post Office Address (if different)

Full name of fourth inventor

Ervin Epstein, Jr.

Inventor's signature

Date

**Residence**

553A Miner Road, Orinda, CA 94563

Citizenship

USA

Post Office Address (if different)

08916140-08349  
45 "T380" 04T980